

# Physio-metabolic and clinical consequences of wearing face masks -Systematic review with meta-analysis and comprehensive evaluation

**Kai Kisielinski** (✉ [kaikisielinski@yahoo.de](mailto:kaikisielinski@yahoo.de))

Orthopaedic and Trauma surgery, Clinical medicine, Private practice, 40212 Düsseldorf, Germany  
<https://orcid.org/0000-0002-2612-6256>

**Oliver Hirsch**

Department of Psychology, FOM University of Applied Sciences, 57078 Siegen, Germany

**Susanne Wagner**

Veterinary medicine, Wagner MSL Management, 15831 Mahlow, Germany

**Barbara Wojtasik**

Department of Genetics and Biosystematics, Faculty of Biology, University of Gdańsk, 80-308, Gdansk, Poland

**Stefan Funken**

Internal medicine, Clinical medicine, Private practice, 47447 Moers, Germany

**Bernd Klosterhalfen**

Institute of Pathology, Dueren Hospital, 52351 Dueren, Germany

**Soumen Kanti Manna**

Biophysics and Structural Genomics Division, Saha Institute of Nuclear Physics, Kolkata 700064, India

**Andreas Prescher**

Institute of Molecular and Cellular Anatomy (MOCA), RWTH Aachen University, 52074 Aachen, Germany

**Pritam Sukul** (✉ [pritam.sukul@uni-rostock.de](mailto:pritam.sukul@uni-rostock.de))

Rostock Medical Breath Research Analytics and Technologies (ROMBAT), Dept. of Anesthesiology and Intensive Care, University Medicine Rostock, 18057 Rostock, Germany <https://orcid.org/0000-0001-5114-1776>

**Andreas Sönnichsen**

Internal medicine, Clinical medicine, Private practice, Gesundheit für Österreich e.V. (Health for Austria), 1030 Vienna, Austria

---

## Systematic Review

**Keywords:** face masks, N95 face mask, surgical mask, risk, adverse effects, long-term adverse effects, health risk assessment, hypercapnia, hypoxia, MIES syndrome

**DOI:** <https://doi.org/10.21203/rs.3.rs-2394501/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Physio-metabolic and clinical consequences of wearing face masks -Systematic review with meta-analysis and comprehensive evaluation

Kai Kisielinski<sup>1,\*</sup>, Oliver Hirsch<sup>2</sup>, Susanne Wagner<sup>3</sup>, Barbara Wojtasik<sup>4</sup>, Stefan Funken<sup>5</sup>, Bernd Klosterhalfen<sup>6</sup>, Soumen Kanti Manna<sup>7</sup>, Andreas Prescher<sup>8</sup>, Pritam Sukul<sup>9,\*</sup>, Andreas Sönnichsen<sup>10</sup>

<sup>1\*</sup>Orthopaedic and Trauma surgery, Clinical medicine, Private practice, 40212 Düsseldorf, Germany

<sup>2</sup>Department of Psychology, FOM University of Applied Sciences, 57078 Siegen, Germany

<sup>3</sup>Veterinary medicine, Wagner MSL Management, 15831 Mahlow, Germany

<sup>4</sup>Department of Genetics and Biosystematics, Faculty of Biology, University of Gdańsk, 80-308, Gdansk, Poland

<sup>5</sup>Internal medicine, Clinical medicine, Private practice, 47447 Moers, Germany

<sup>6</sup>Institute of Pathology, Dueren Hospital, 52351 Dueren, Germany

<sup>7</sup>Biophysics and Structural Genomics Division, Saha Institute of Nuclear Physics, Kolkata 700064, India

<sup>8</sup>Institute of Molecular and Cellular Anatomy (MOCA), RWTH Aachen University, 52074 Aachen, Germany

<sup>9\*</sup>Rostock Medical Breath Research Analytics and Technologies (ROMBAT), Dept. of Anesthesiology and Intensive Care, University Medicine Rostock, 18057 Rostock, Germany

<sup>10</sup>Internal medicine, Clinical medicine, Private practice, Gesundheit für Österreich e.V. (Health for Austria), 1030 Vienna, Austria

\*Corresponding authors: [kaikisielinski@yahoo.de](mailto:kaikisielinski@yahoo.de) & [pritam.sukul@uni-rostock.de](mailto:pritam.sukul@uni-rostock.de)

## Abstract

**Background:** As face masks are a mandatory public health intervention during the COVID-19 pandemic, adverse effects require substantiated investigation.

**Methods:** A systematic review of 2168 studies yielded 54 publications for synthesis and 37 studies for meta-analysis (on  $n=8641$ ,  $m=2482$ ,  $f=6159$ ,  $age=34.8\pm 12.5$ ). The median trial duration was only 18 min (IQR=50) for our comprehensive evaluation of mask induced physio-metabolic and clinical outcomes.

**Results:** We found significant effects in both medical masks with a greater impact regarding the N95. These effects included decreased SpO<sub>2</sub> (overall SMD=-0.24, 95%CI=-0.38 to -0.11,  $p=0.0004$ ) and minute ventilation (SMD=-0.72, 95%CI=-0.99 to -0.46,  $p<0.00001$ ), simultaneously increased blood-CO<sub>2</sub> (SMD=+0.64, 95%CI=0.31–0.96,  $p=0.0001$ ), heart rate (N95: SMD=+0.22, 95%CI=0.03–0.41,  $p=0.02$ ), systolic blood pressure (surgical: SMD=+0.21, 95%CI=0.03–0.39,  $p=0.02$ ), skin temperature (overall SMD=+0.80 95%CI 0.23–1.38,  $p=0.006$ ) and humidity (SMD +2.24, 95%CI=1.32–3.17,  $p<0.00001$ ). Effects on exertion (overall SMD=+0.9, surgical=+0.63, N95=+1.19), discomfort (SMD=+1.16), dyspnoea (SMD=+1.46), heat (SMD=+0.70) and humidity (SMD=+0.9) were significant in 373 cases with a robust relationship to mask wearing ( $p<0.006$  to  $p<0.00001$ ). Pooled symptom prevalence was significant in users ( $n=8128$ ) for: headache (62%,  $p<0.00001$ ), acne (38%,  $p<0.00001$ ), skin irritation (36%,  $p<0.00001$ ), dyspnoea (33%,  $p<0.00001$ ), heat (26%,  $p<0.00001$ ), itching (26%,  $p<0.00001$ ), voice disorder (23%,  $p<0.03$ ) and dizziness (5%,  $p=0.01$ ).

**Discussion:** Masks interfered with O<sub>2</sub>-uptake and CO<sub>2</sub>-release and compromised respiratory compensation. Though evaluated wearing durations do not represent daily/prolonged use, outcomes independently validate mask-induced exhaustion-syndrome (MIES). MIES can have long-term clinical consequences, especially for vulnerable groups.

**Conclusion:** Face mask side-effects must be assessed (risk-benefit) against the available evidence of their effectiveness against viral transmissions.

## Introduction

The use of face masks has been restricted to professionals for decades. In the health-care setting, masks constituted a mandatory self-protective and third-party protective measure for medical personnel prior to COVID-19 pandemic and there is no doubt about the efficacy of masks in reducing transmission of pathogens, especially bacteria. In 2020, many scientists and leaders started to believe that the use of masks could also provide protection against viral transmission, although evidence for the effectiveness of this measure was only weak<sup>1</sup>. In the meantime, a large number of publications on this topic cannot be overlooked<sup>2,3</sup>.

During the 2019 SARS-CoV-2 outbreak face masks were deployed as a mandatory public health measure for the general population in many countries around the world, making them one of the most important universal life-style attributes that directly affects how we breathe. As with any other preventive measure and/or intervention, masks also have specific advantages and disadvantages. While certain properties may have justified their invention and application in the past, e.g. retention of bacteria during surgical wound care and operations, at present the question needs to be addressed as to the long-term effects widespread mask wearing may have on normal breathing. It is noteworthy that the compulsory wearing of masks for the entire population provided good research conditions and consequently numerous publications dealing with the adverse effects of mask wearing<sup>4-11</sup>. Various volatile metabolites are produced from *in vivo* biochemical and metabolic pathways and their concentrations in exhaled breath provide immediate physiological<sup>12,13</sup>, metabolic<sup>14,15</sup> and pathological<sup>16,17</sup> magnitudes with the possibility of monitoring various processes and interventions including therapies<sup>18,19</sup>. A recent observational study reported continuous respiratory and haemodynamic changes along with corresponding alteration in exhaled volatile metabolites (viz. potentially originate at the cellular/organ levels and via microbial metabolic processes) and has raised significant concerns upon the immediate, progressive, transient and long-term side-effects of FFP2/N95 and surgical masks in adults (aged between 20 – 80 years) at rest<sup>20</sup>. Previously, based on other numerous publications major concerns were raised in a large-scale scoping review<sup>8</sup>. Furthermore, this recent scoping review on mask driven adverse effects and health risks has summoned for a systematic review.

Though some important systematic reviews regarding masks and their effects already exist<sup>21-24</sup>, they are predominantly restricted to healthy and sportive individuals<sup>21,23</sup>. Due to the exclusion of children, pregnant women and diseased patients from these evaluations and conclusions<sup>22,25</sup>, the reviews do not provide sufficient evidence that masks can be employed in the general population as safe protective measures. Moreover, the application of fixed statistical models<sup>21</sup>, use of narratives rather than quantitative analysis and statistics (despite claiming to be systematic)<sup>26</sup>, focus only on health care workers and their complaints<sup>25</sup>, as well as comparing the different mask types without any baseline/control group<sup>25</sup> were ubiquitous limitations of those studies. Physiological systematic reviews based purely on physiological effects of masks limit data interpretations to normal physio-metabolic fluctuations i.e. beyond the domain of pathophysiological compensatory mechanisms (especially in the elderly and those with diminished compensatory reserves) and/or acute/chronic subliminal changes in the human microbiome<sup>22,24</sup>. Similarly, other manuscripts do not address subjective parameters, prevalence of symptoms and discomfort during mask use and concomitant physical changes such as heat and temperature in detail<sup>21,23</sup>. Therefore, the systematic reviews available to date neither address possible symptoms of mask use for the general population nor their exact prevalence. In addition, the transferability of the outcomes of said systematic reviews to the general population is very limited and they do not fulfil the actual requirements of clinical and inclusive evaluation, especially from the views and perspective of physicians and clinicians. Including young, old, healthy and ill people for the systematic analysis of physiological, metabolic and clinical data could complete the possible comprehensive impact of mask-wearing on the general

population. In contrast to the above-indicated studies, our systematic review is aimed to quantify the biochemical/metabolic, physical, physiological changes along with the appearance of subjective and clinical symptoms in face mask users and analyse them from a clinician's and physician's holistic perspective.

## **Materials & Methods**

### *Registration*

This meta-analysis was registered with the international prospective register of systematic reviews (PROSPERO) under the record CRD42021256694 at the National Institute for Health Research (NIHR) and performed in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement <sup>27</sup>.

### *Inclusion and exclusion criteria*

The aim was to study adverse effects of face masks on metabolic, physiological, physical, psychological and individualised parameters. The use of cloth masks, surgical masks and N95/FFP-2 masks were the intervention of interest. Humans of all ages and genders, who were evaluated in controlled intervention studies and observational studies have been included in our comprehensive evaluation. Case reports, narrative reviews, case series and expert opinions were excluded. The main outcomes considered were peripheral oxygen saturation (SpO<sub>2</sub>), carbon dioxide levels in blood, temperature, humidity, heart rate, respiratory rate, tidal volume and minute ventilation, blood pressure, exertion, dyspnoea, discomfort, headache, skin changes, itching, psychological stress and symptoms during the use of face masks.

### *Literature retrieval strategy*

First, a comprehensive search term was developed. Then, PubMed, Embase and Cochrane Library databases were searched. The search was performed until 31st December 2021. There were no restrictions in publication date. Literature that was neither English nor German language was excluded. Additionally, forward-looking data (e.g. available as preprint, but not published in a peer-reviewed scientific journal at time of completing this meta-analysis) was considered for discussion, but not included in the meta-analysis.

### *Literature screening and data extraction*

Search terms were created according to the criteria defined in the PICO scheme <sup>28</sup>. The specific search terms were: (face mask\* [tw], FFP1 [tw] FFP2 [tw], FFP3 [tw], N99 [tw], N97 [tw], N95 [tw], respiratory protective device\* [tw], air-purifying respirator\* [tw], surgical mask\* [tw]) and (risk\* or adverse effect\* [tw], adverse event\* [tw], side effect\* [tw], psycho\* [tw], hypoxia [tw], hypercapnia [tw], headache [tw], dead space [tw], safety [tw], carbon dioxide [tw]), not infants, not neonatal, not newborn, not endoscopy, not CPAP, not intubate\*, not propofol, not resuscitation, not mechanical ventilation [tw], not foetus. The asterisk in the search algorithm here '\*' stands for the extension of the spelling with different possible letter combinations (e.g. face mask\* with \*= s, or \*=ed, or \*=ing). The abbreviation '[tw]' stands for title word.

The retrieved titles and abstracts were then screened and assessed for predefined inclusion criteria by at least three authors. Study design, methodology, interventions, primary and secondary

outcomes and language were evaluated using the web-based program Rayyan — a web and mobile app for systematic reviews<sup>29</sup>. Full texts of all potentially relevant articles were independently assessed for inclusion by two authors. Full-text exclusions and reasons have been documented. Data of included full texts were extracted: Author and year, type of study, aim of the study, intervention/control, sample size, follow-up, outcomes, funding, setting/country, age, sex, comorbidities, medications, functional status and cognitive status of participants, results, main findings, and limitations. Descriptive data was extracted by one author and checked by a senior author. If discrepancies occurred or authors disagreed, a senior author was involved in and a consensus was found<sup>30</sup>.

#### *Risk of bias assessment of the included studies*

The quality assessments were carried out using various tools, depending on the type of study. If systematic reviews and meta-analyses were included, these were assessed using the AMSTAR-2 checklist<sup>31</sup>. Interventional studies were examined using the manual “Assessment of the risk of bias in clinical studies” from the Cochrane Collaboration (Cochrane RoB-2)<sup>32</sup>. Observational studies were checked with the CASP (Critical Appraisal Skills Program) using standardised forms<sup>33</sup>.

#### *Statistical analysis*

A meta-analysis was carried out, if enough studies with the same research question were found among the randomised, non-randomised controlled trials and observational studies. A subgroup analysis was conducted, where possible, for different mask types (N95/surgical) and even compared the mask types with each other (N95 vs surgical mask). The program "RevMan-5.4.1", which was developed for Cochrane Reviews was used. As we anticipated a considerable between-study heterogeneity -the random effects model was used to pool effect sizes<sup>34</sup>. The results were graphically depicted in forest plots. Subgroup analyses were performed and a Q test was calculated to examine significant subgroup differences. Study heterogeneity was assessed using Cochrane's Q test, T2 according to DerSimonian / Laird<sup>35</sup>, and I<sup>2</sup> according to Higgins / Thompson<sup>36</sup>. Where possible, a funnel plot was created to investigate publication bias. If this showed an abnormal result and there were at least ten studies evaluating the same question, Egger's test<sup>37</sup> was carried out. For the analysis of metabolic and physiological changes all controlled intervention studies in which measurements were taken during physical activity with face masks were included. We excluded resting conditions since these are not representative for real life settings and pre-post studies to ensure study-comparability. In addition, by excluding rest situations of the mostly healthy study participants, our approach was able to represent the possible effects better in elderly and ill individuals (e.g. with compromised compensation mechanisms), all of whom are a significant part of the general population. This also helped to reduce heterogeneity (I<sup>2</sup>). Neither for the results of the systolic blood pressure (SBP) nor the temperature did we follow this approach. Studies in which measurements were taken during rest and moderate physical activity were included in the meta-analysis of the physical outcome on SBP to obtain an evaluable number of studies and to ensure a better comparability and lower heterogeneity (exclusion of heavy load exercise conditions). In order to gather more available data for evaluating the temperature, we included two pre-post studies containing a resting condition using valid methodology and exact temperature measurements. This clearly reduced the heterogeneity index I<sup>2</sup>. For the meta-analysis of the resultant CO<sub>2</sub>-blood-content the joint evaluation of different experimental CO<sub>2</sub> measurements (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>) in mmHg was justified by the following facts:

- 1) “ETCO<sub>2</sub> and PtCO<sub>2</sub> measurements both provide an estimation of PaCO<sub>2</sub>”<sup>38</sup>.
- 2) "End-tidal CO<sub>2</sub> (ETCO<sub>2</sub>) has been considered as a reliable estimate of arterial PCO<sub>2</sub>, in healthy subjects"<sup>39</sup>.

- 3) "PtCO<sub>2</sub> reliably reflects PaCO<sub>2</sub>, irrespective of sensor location" <sup>40</sup>.
- 4) "Transcutaneous CO<sub>2</sub> (PtCO<sub>2</sub>) devices provide another option for the continuous non-invasive estimation of PaCO<sub>2</sub>, overcoming the limitations posed by end-tidal CO<sub>2</sub> analysis" <sup>39</sup>.
- 5) "ETCO<sub>2</sub> monitoring tends to underestimate PaCO<sub>2</sub> levels" <sup>38</sup>.

For meta-analysis of measured sensations, all studies in which measurements were mainly taken during physical activity were included. This helped to ensure comparability, lower heterogeneity and the above mentioned aims to draw conclusions on the general population under conditions resembling real life settings. However, an exemption was made for the sensation 'discomfort': To allow evaluable study numbers, we included one pre-post study with resting condition, however, with valid methodology and exact discomfort evaluations <sup>41</sup>. Even if this study had not been included, the result would be significant and unambiguous, however with a slightly larger 95% CI. Our systematic review also referenced studies aiming to assess the prevalence of sensations and symptoms under mask use. Therefore, we conducted an additional meta-analysis of these observational studies to document the pooled prevalence in mask use. Prevalence was calculated as total number of symptoms per 100 mask wearers. In studies where the standard error (SE) was not reported, we calculated it from the prevalence using the following formula:  $SE = \sqrt{p(1-p)/n}$  with a 95% CI =  $p \pm 1.96 \times SE$ ; where,  $p$  = Prevalence. This statistical approach to quantify a pooled prevalence from observational studies has been previously reported <sup>34</sup>. Meta-analysis was performed using RevMan (Version 5.4.1). The heterogeneity of each meta-analysis was assessed and then the random effects model was used to calculate the pooled prevalence. We conducted subgroup analysis where possible for mask type (N95/surgical). Funnel plots were used to study the possibility of publication bias as described above.

The inclusion of observational studies, particularly for the prevalence analysis in our meta-analysis is justified because these are particularly suitable to investigate exposures that are difficult or impossible to investigate in randomised controlled trials (RCTs), e.g. air pollution or smoking. In addition, observational studies are important to investigate causes with a long latency period, such as carcinogenic effects of environmental exposures or drugs <sup>42</sup>. Thus, possible adverse long-term effects of masks, i.e. comparable to the environmental hazards, appeared to be particularly detectable through observational studies.

Finally, the random statistical control calculations of our results were performed for quality assurance via the R software (R Foundation for Statistical Computing, Vienna, Austria, version 4.0.1) and packages metafor, dmetar, meta <sup>30</sup>. Knapp-Hartung adjustments to control for the uncertainty in the estimate of the between-study heterogeneity were used in these calculations which are controversial as they result in wider confidence intervals and are also suspected to be anti-conservative even though the effects are very homogeneous <sup>30</sup>.

## Results

### General findings

#### Literature characteristics

Of the 2168 screened records, 54 studies were included for qualitative analysis (see extraction tables, Table 1) and 37 for statistical meta-analysis (Figure 1). Among the 54 studies, 23 were intervention studies, and 31 were observational studies. The 23 intervention studies consisted of 14 randomised controlled trials (RCT's) and 9 non-randomised controlled trials (nRCT's). Of the 31 observational studies, 17 works raised measured values, and 14 were questionnaire studies.

**Table 1 A-C:** Overview of 54 included studies. A randomised controlled trials, B non-randomised controlled trials and C observational studies

**Table 1A: Included 14 randomised controlled trials**

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
<b>Bertoli 2020</b>	Randomized, two-period cross-over self-control trial	Wearing N95 respirator vs no facemask during indirect calorimetry	N=10	5 min	oxygen consumption (VO <sub>2</sub> ), carbon dioxide production (VCO <sub>2</sub> ), Resting Energy Expenditure (REE)
<b>Butz 2004</b>	Blinded, randomized cross over study	Wearing two types of surgical masks vs no mask	N=15	30 min	CO <sub>2</sub> under masks, PtCO <sub>2</sub> (partial transcutaneous CO <sub>2</sub> pressure) while wearing masks for 30 min, HR, RR (respiratory rate), SpO <sub>2</sub>
<b>Dirol 2021</b>	Prospective randomized cross-over study	Six-minute walking test (6MWT) with and without surgical mask. Mask-discomfort questionnaire was applied before and after 6 MWT with the mask	N=100	6 min	RR, HR, SpO <sub>2</sub> , EtCO <sub>2</sub> , discomfort questionnaire
<b>Fikenzer 2020</b>	Prospective cross-over study	Wearing no mask (nm) vs surgical mask (sm) vs FFP2/N95 mask (ffpm), cardiopulmonary and metabolic responses monitored by ergo-spirometry and impedance cardiography	N=12	10 min	FVC (forced vital capacity), FEV1 (forced expiratory volume in 1 s), Tiffenau index, peak expiratory flow (PEF), HR, stroke volume, cardiac output, arterio-venous oxygen content difference, systolic blood pressure (SBP), diastolic blood pressure (DBP), ventilation in liters/minute (VE), RR, tidal volume (VT), pH, partial pressure of carbon dioxide (PaCO <sub>2</sub> ), partial pressure of oxygen (PaO <sub>2</sub> ), lactate Pmax, Pmax/kg, VO <sub>2</sub> max/kg, heart rate recovery (HRR): HRR-1 min, HRR-5 min. Discomforts (VAS): humid, hot, breath resistance, itchy, tight, salty, unfit, odor, fatigue, overall discomfort.
<b>Georgi 2020</b>	Prospective randomized cross-over study	wearing no mask (nm) vs community vs surgical mask (sm) vs FFP2/N95 mask (ffp treadmill: baseline, 50 W, 75W, 100W)	N=24	9 min	HR, RR, SBP, DBP, PtCO <sub>2</sub> , SpO <sub>2</sub> , main symptoms questionnaire
<b>Goh 2019</b>	Randomized, two-period cross-over	Wearing N95 respirator vs wearing N95 respirator with microfan	N=106	15 min	EtCO <sub>2</sub> , comfort level with

*Consequences of wearing face masks*

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
<b>Hua 2020</b>	self-control trial Prospective randomized crossover trial	vs wearing no facemask during common physical activities Two and 4 hours after donning the masks, adverse reactions and perceived discomfort and noncompliance were measured.	N=20	240 min	visual analogue scale (VAS) Skin parameters: Skin hydration, transepidermal water loss, erythema, pH and sebum secretion
<b>Kim J.H. 2013</b>	Randomized, self-control trial	Wearing N95 respirator (partly with exhalation valve) vs wearing no facemask (NM) during a low-moderate work-rate (5.6km/h)	N=20	60 min	HR, RR, transcutaneous carbon dioxide, SpO <sub>2</sub>
<b>Kim J.H. 2015</b>	Randomized, two-period controlled trial	Wearing N95 respirator and no mask during one hour of mixed sedentary activity and moderate exercise during pregnancy vs non pregnant women	N= 16 vs 16	60 min	SBP, DBP, mean arterial pressure, HR, stroke volume, cardiac output, total peripheral resistance, RPE, SpO <sub>2</sub> , PtCO <sub>2</sub>
<b>Kim J.H. 2016</b>	Randomized, self-control trial	Wearing N95 respirator vs wearing P100 respirator vs wearing no mask during 1 hour of treadmill exercise (5.6 km/h) in an environmental chamber (35°C, relative humidity 50%)	N=12	60 min	Fit factor, rectal temperature, mean skin temperature, facial skin temperature under respirator, SpO <sub>2</sub> , PtCO <sub>2</sub> , HR, RR, breathing comfort, thermal sensation, exertion (Borg scale)
<b>Mapelli 2021</b>	interventional, prospective, randomized, double-blind and cross-over study	Wearing no mask surgical mask or N95 mask and performing consecutive cardiopulmonary exercise tests (CPETs) at least 24 hours apart but within 2 weeks	N=12	10 min	Ventilation (VE), Oxygen intake VO <sub>2</sub> , VCO <sub>2</sub> production, respiratory gases,: expiratory O <sub>2</sub> (ETO <sub>2</sub> ) and expiratory CO <sub>2</sub> (ETCO <sub>2</sub> ), Heart rate (HR), hemoglobin saturation (SaO <sub>2</sub> ), blood pressure (DBP and SBD), dyspnea (Borg scale), Spirometry, Maximal Inspiratory pressure (MIP) and Maximal Expiratory Pressure (MEP)
<b>Roberge 2014</b>	Randomized, two-period controlled trial	Wearing an N95 FFR during exercise and postural sedentary activities over a 1-hour period on pregnant women vs control	N= 22/22	60 min	Core temperature, cheek temperature, abdominal temperature, HR, RR, RPE, perceived heat (RHP)
<b>Wong A.Y.-Y 2020</b>	Randomized, two-period self-controlled trial	Wearing a facemask vs not wearing a facemask during graded treadmill (10% slope) walking at 4 km/h for 6 min	N=23	6 min	HR, RPE
<b>Zhang 2021</b>	Prospective randomized cross-over study	Exercises (cycle ergometer) with and without surgical masks (mask-on and mask-off) were analyzed	N=71	8 min	test duration, maximum power, RPE score, Borg dyspnea scale, Oxygen consumption (V. O <sub>2</sub> ), carbon dioxide production (V.CO <sub>2</sub> ), metabolic equivalent (MET), respiratory exchange

Consequences of wearing face masks

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
					rate (RER), and percentage of oxygen uptake at anaerobic threshold (AT) in predicted maximal oxygen uptake, inspiratory time (Ti), expiratory time (Te), RR, VT, VE, end-tidal oxygen partial pressure (EtO <sub>2</sub> ), EtCO <sub>2</sub> , oxygen ventilation equivalent (VE/V.O <sub>2</sub> ), and carbondioxide equivalent (V <sub>E</sub> /VCO <sub>2</sub> )
<b>Legend:</b>					
<p><i>AT, anaerobic threshold; DBP = diastolic blood pressure; EtCO<sub>2</sub> = end-tidal CO<sub>2</sub> partial pressure; ESRD = end stage renal disease; TEWL= trans-epidermal water loss; FEV1 = forced expiratory volume in 1 sec; FVC = forced vital capacity; HCW = health care worker; HD=haemodialysis; HR = heart rate; MEP = maximal expiratory pressure, MET = metabolic equivalent; MIP =maximal inspiratory pressure; PEF = peak expiratory flow; PetCO<sub>2</sub> = end-tidal carbon dioxide pressure; PetO<sub>2</sub> =end-tidal oxygen pressure ; PI = perfusion index; PPE = personal protective equipment; PtCo<sub>2</sub> = partial transcutaneous CO<sub>2</sub> pressure; RER = respiratory exchange ratio; RPE = rated perceived exertion; RR = respiratory rate; RR =respiratory rate; SaO<sub>2</sub> =hemoglobin oxygen saturation; SBP = systolic blood pressure; SpO<sub>2</sub> = oxygen saturation; Te= expiratory time; Ti = inspiratory time; Tiot =Inspiratory + expiratory time; TV = tidal volume; V CO<sub>2</sub> =carbon dioxide production; V O<sub>2</sub> = oxygen uptake; VE = ventilation in liters/min; VE = ventilation; VT = tidal volume,</i></p>					

**Table 1B: Included 9 non-randomised controlled trials**

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
<b>Bharatendu 2020</b>	Cross-sectional self-control trial	Wearing N95 respirator vs no facemask	N=154	5 min	Mean flow velocity (MFV), Pulsatility index, end-tidal carbon dioxide partial pressure (EtCO <sub>2</sub> )
<b>Coniam 2005</b>	Two-period controlled trial	Wearing surgical masks (WM) vs no facemask (NM) during oral examination	N=186	10 min	Pronunciation, vocabulary, grammar, comprehensibility, audibility
<b>Epstein 2020</b>	Multiple cross-over, self-control trial	Wearing N95 respirator vs wearing surgical mask vs no facemask during maximal exercise test	N16	18 min	HR, RR, SpO <sub>2</sub> , rated perceived exertion (RPE), end-tidal carbon dioxide (EtCO <sub>2</sub> )
<b>Lee 2011</b>	Two-period self-controlled trial	Wearing N95 respirator vs no facemask during rhinomanometry	N=14	30 sec	Inspiration breathing resistance increment, expiration breathing resistance increment, breathing volume decrement
<b>Roberge 2010</b>	Multiple cross-over, self-control trial	Wearing an N95 FFR vs N95 FFR with exhalation valve vs no mask during 1-hour treadmill walking sessions, at 1.7 miles/h and at 2.5 miles/h	N=10	60 min	FFR dead space gases, CO <sub>2</sub> saturation, O <sub>2</sub> saturation, RR, VT, VE, HR
<b>Roberge 2012</b>	Two-period self-control trial	Wearing a surgical mask for 1 hour during treadmill exercise at 5.6 km/h vs the same exercise with no mask	N=20	60 min	Core temperature, cheek temperature, abdominal

## Consequences of wearing face masks

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
					temperature, HR, RR, RPE, Perceived heat (RHP)
<b>Scarano 2020</b>	Two-period self-controlled trial	Wearing a surgical mask for 1 hour vs wearing N95 respirator for 1 hour vs baseline	N=20	60 min	Humidity, heat, breathing difficulty, discomfort, mask touching, perioral temperature
<b>Shenal 2012</b>	Multiple cross-over self-controlled field trial	Wearing one of seven respirators or medical mask during an 8-hour working period vs no mask	N=27	480 min	Discomfort, RPE
<b>Tong 2015</b>	Two-period self-controlled trial	Breathing through N95 mask materials during rest and exercise of predetermined intensity vs breathing ambient air	N=19	50 min	Oxygen consumption (VO <sub>2</sub> ), carbon dioxide production (VCO <sub>2</sub> ), VT, RR, VE, expired oxygen (FeO <sub>2</sub> ), expired carbon dioxide (FeCO <sub>2</sub> ), inspired oxygen (FiO <sub>2</sub> ), inspired carbon dioxide (FiCO <sub>2</sub> )
<b>Legend:</b>					
<p><i>AT, anaerobic threshold; DBP = diastolic blood pressure; EtCO<sub>2</sub> = end-tidal CO<sub>2</sub> partial pressure; ESRD = end stage renal disease; TEWL = trans-epidermal water loss; FEV<sub>1</sub> = forced expiratory volume in 1 sec; FVC = forced vital capacity; HCW = health care worker; HD=haemodialysis; HR = heart rate; MEP = maximal expiratory pressure, TMET1 = metabolic equivalent; MIP =maximal inspiratory pressure; PEF = peak expiratory flow; PetCO<sub>2</sub> = end-tidal carbon dioxide pressure; PetO<sub>2</sub> =end-tidal oxygen pressure ; PI = perfusion index; PPE = personal protective equipment; PtCo<sub>2</sub> = partial transcutaneous CO<sub>2</sub> pressure; RER = respiratory exchange ratio; RPE = rated perceived exertion; RR = respiratory rate; RR =respiratory rate; SaO<sub>2</sub> =hemoglobin oxygen saturation; SBP = systolic blood pressure; SpO<sub>2</sub> = oxygen saturation; Te= expiratory time; Ti = inspiratory time; Ttot =Inspiratory + expiratory time; TV = tidal volume; V'CO<sub>2</sub> =carbon dioxide production; V'O<sub>2</sub> = oxygen uptake; VE = ventilation in liters/min; VE = ventilation; VT = tidal volume,</i></p>					

**Table 1C: Included 31 observational studies**

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
<b>Beder 2008</b>	Longitudinal and prospective observational study	Wearing surgical mask during major operations vs baseline	N=53	60-240 min	SpO <sub>2</sub> ,(oxygen saturation) HR (heart rate)
<b>Choudhury 2020</b>	Prospective cohort study	Wearing N95 respirator during light work vs wearing full PPE during heavy work vs baseline	N=75	240 min	HR,SpO <sub>2</sub> , Perfusion Index (PI), RPE (rated perceived exertion), modified Borg scale for dyspnoea
<b>Foo 2006</b>	Survey study	Self-administered questionnaire healthcare workers	N=322	480 min	Prevalence of adverse skin reactions
<b>Forgie 2009</b>	Cross-sectional survey study	Self-administered questionnaire	N=80	Not given	Mask/Shield preference Mask results, Shield resul
<b>Heider 2020</b>	Cross-sectional survey study	Validated Voice Handicap Index (VHI)-10 questionnaire and self administered questionnaire	N=221	480 min	Vocal symptoms, Spanish validated Voice Handicap Index (VHI)-10 questionnaire
<b>Islam 2022</b>	Prospective cross-over self-control	Wearing FFP2 (N95) mask for 30 mins under sitting condition in	N = 10	30 min	Saha Institute of Nuclear

*Consequences of wearing face masks*

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
	study	an air-conditioned room			Physics, Department of Atomic Energy, Government of India
<b>Jafari 2021</b>	Cross-sectional study	Self-administered questionnaire, SpO <sub>2</sub> , HR and venous blood samples	N=243	240 min	RR, HR, SpO <sub>2</sub> , salivary metabolic signature
<b>Kao 2004</b>	Prospective observational study	Wearing N95 respirator during haemodialysis vs baseline	N=39	240 min	HR, RR, systolic blood pressure (SBP), diastolic blood pressure (DBP), PaO <sub>2</sub> , PaCO <sub>2</sub> discomfort rates
<b>Klimek 2020</b>	Cross-sectional Survey study	Visual Analogue Scales (VAS) to document patient-reported symptoms and diagnostic findings	N=46	120 min	Visual Analogue Scales (VAS) to document patient-reported symptoms of: rhinitis, rhinorrhea. Mucosal irritation, secretion and edema in nasal endoscopy was graded
<b>Kyung 2020</b>	Prospective panel study	Wearing N95 respirator during 6 minute walking test vs baseline	N=97	6 min	SBP, DBP, HR, RR, EtCO <sub>2</sub> , SpO <sub>2</sub> ,
<b>Lan 2020</b>	Cross-sectional Survey study	Self-administered questionnaire	N=542	360 min	Prevalence of adverse skin reactions
<b>Li 2005</b>	Prospective observational study	Exercise on a treadmill while wearing the protective facemasks	N=10	100 min	HR, temperature and humidity (outside and inside the facemask), SBP, DBP, mask outer humidity, face microclimate humidity, chest microclimate humidity, mask outside temperature, face microclimate temperature, face skin temperature, chest microclimate temperature, subjective sensations: humiditty, heat, breath resistance, itching, tightness, feeling salty, feeling unfit, feeling odorous, fatigue, overall discomfort
<b>Lim 2006</b>	Survey study	Self-administered questionnaire	N=212	240 min	Prevalence of headaches
<b>Luckman 2020</b>	survey study using online experimental setting	Self-administered questionnaire and experimental online setting	N=400	Not given	Risk compensation with reduced physical distancing (standing, sitting, walking)
<b>Matusiak 2020</b>	Cross-sectional	Self-administered questionnaire	N=876	Not	Difficulty in breathing,

*Consequences of wearing face masks*

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
	Survey study			given	warming/sweating glasses misting up, slurred speech, itch
<b>Mo 2020</b>	Retrospective observation cross over cohort study	Wearing surgical mask vs not wearing: compare to former hospitalisations. Including criteria: Patients who were hospitalized three or more times and at least two times before mask mandates	N=23	7 min	Vital signs: temperature, HR, RR, SBP, DBP, serum and blood gas analysis, inpatient days (days). clinical parameters, including ion concentration of serum, vital signs, inflammation markers and artery blood gas.
<b>Naylor 2020</b>	Survey study	Self-administered online questionnaire.	N=129	Not given	Effects of certain aspects of lockdown, including face masks, social distancing, and video calling, on participants behavior, emotions, hearing performance, practical issues, and tinnitus.
<b>Ong 2020</b>	Cross-sectional survey study	Self-administered questionnaire.	N=158	360 min	PPE usage patterns, occupation, underlying comorbidities
<b>Park 2020</b>	Prospective cohort study	Wearing KF94 respirator for 6 hours vs baseline	N=21	360 min	Skin temperature increase, skin redness, skin hydration, sebum level, skin elasticity, trans-epidermal water loss
<b>Pifarre 2020</b>	Prospective trial	No mask baseline vs. Mask baseline. Subjects wearing a mask immediately after a 21-flex test performed the Ruffier protocol	N=8	5-7 min	PaO <sub>2</sub> , PaCO <sub>2</sub> , SpO <sub>2</sub> , HR
<b>Prousa 2020</b>	Cross-sectional survey study	Self-administered questionnaire	N=1010	Not given	Wearing time, discomfort Stress, Tricks, psychovegetative complaints, positive feelings, aggression, depression
<b>Ramirez-Moreno 2020</b>	Cross-sectional study in healthcare workers	Self-administered questionnaire	N=306	420 min	Work type, type of face mask, number of hours worn per day (SD). pre-existing headache, comorbidity, other symptoms, Sleep disturbance, loss of concentration, irritability, photophobia, sonophobia, sickness/vomiting
<b>Rebmann 2013</b>	Multiple cross-over, self-control trial	Wearing only an N95 or an N95 with mask overlay for a 12-hour	N=10	720 h	SBP, DBP, CO <sub>2</sub> saturation,

Consequences of wearing face masks

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
		shift vs baseline			SpO <sub>2</sub> , HR, headache, nausea, light-headedness, visual challenge
<b>Rosner 2020</b>	Cross-sectional study in healthcare workers	Self-administered questionnaire	N=343	360 min	Acne, headache, skin breakdown (nose bridge, cheeks, chin. behind ears), impaired cognition
<b>Sukul 2022</b>	Two-period controlled trial	Wearing a surgical or N95 mask during rest (young to mid-aged adults were measured for 30 min and older adults were measured for 15 min)	N=30	15-30 min	Exhaled breath profiles within mask space by high-resolution real-time mass-spectrometry (PTR-ToF-MS): aldehydes, hemiterpene, organosulfur, short-chain fatty acids, alcohols, ketone, aromatics, nitrile and monoterpene. Haemodynamic parameters: SpO <sub>2</sub> , PETCO <sub>2</sub> , HR, RR, SBP, DBP, cardiac output, exhaled oxygen, humidity.
<b>Szczesniak 2020</b>	Survey study	Self-administered online questionnaire After mask restrictions vs before mask restrictions	N=1476 vs 564	Not given	Employment status, place of residence, worktime per week, somatic symptoms, anxiety and insomnia, social dysfunction, depression
<b>Szepietowski 2020</b>	Survey study	Self-administered online questionnaire	N=2307	Not given	itch, mask types used, duration of mask use per day
<b>Techasatian 2020</b>	Prospective cross-sectional survey study	Self-administered questionnaire	N=833	480 min	Factors associated with adverse skin reaction, risk factors for adverse skin reaction, differences between HCW and non-HCW
<b>Thomas 2011</b>	Two-period controlled trial	Comparing the ability to accurately record 20 randomized aviation terms transmitted over the radio by a helicopter emergency medical services (HEMS) pilot wearing a surgical facemask and six different N95s with and without the aircraft engine operating	N=3	Not given	Accurately record 20 terms transmitted over the radio by (HEMS) pilot wearing a surgical facemask or N95 mask
<b>Toprak 2021</b>	Prospective observational study	surgical vs N-95 mask n=149 vs n=148	N=297	35 min	Maternal vital signs: SBP, DBP, HR, RR, fever centigrade, SpO <sub>2</sub>
<b>Tornero-Aguilera 2021</b>	Two-period controlled trial	Wearing a surgical facemask vs not wearing a facemask during 150 min university lessons	N=50	150 min	Mental fatigue perception, reaction time (ms) SpO <sub>2</sub> , mean RR (ms), mean HR

## Consequences of wearing face masks

Author and year	Study design	Intervention/control	Sample size	Time	Outcomes
					(bpm) square root of the mean value of the sum of squared differences of all successive R-R intervals (RMSSD) (ms), low frequency (LF) and high-frequency (HF) normalized units (n.u.), SD1 (ms), SD2 (ms)
<p><b>Legend:</b>  <i>AT</i>, anaerobic threshold; <i>DBP</i> = diastolic blood pressure; <i>EtCO<sub>2</sub></i> = end-tidal CO<sub>2</sub> partial pressure; <i>ESRD</i> = end stage renal disease; <i>TEWL</i>= trans-epidermal water loss; <i>FEV1</i> = forced expiratory volume in 1 sec; <i>FVC</i> = forced vital capacity; <i>HCW</i> = health care worker; <i>HD</i>=haemodialysis; <i>HR</i> = heart rate; <i>MEP</i> = maximal expiratory pressure, <i>TMET1</i> = metabolic equivalent; <i>MIP</i> =maximal inspiratory pressure; <i>PEF</i> = peak expiratory flow; <i>PetCO<sub>2</sub></i> = end-tidal carbon dioxide pressure; <i>PetO<sub>2</sub></i> =end-tidal oxygen pressure ; <i>PI</i> = perfusion index; <i>PPE</i> = personal protective equipment; <i>PtCo<sub>2</sub></i> = partial transcutaneous CO<sub>2</sub> pressure; <i>RER</i> = respiratory exchange ratio; <i>RPE</i> = rated perceived exertion; <i>RR</i> = respiratory rate; <i>RR</i> =respiratory rate; <i>SaO<sub>2</sub></i> =hemoglobin oxygen saturation; <i>SBP</i> = systolic blood pressure; <i>SpO<sub>2</sub></i> = oxygen saturation; <i>Te</i>= expiratory time; <i>Ti</i> = inspiratory time; <i>Ttot</i> =Inspiratory + expiratory time; <i>TV</i> = tidal volume; <i>V̇CO<sub>2</sub></i> =carbon dioxide production; <i>V̇O<sub>2</sub></i> = oxygen uptake; <i>VE</i> = ventilation in liters/min; <i>VE</i> = ventilation; <i>VT</i> = tidal volume,</p>					

### *Quality appraisal*

The quality of the studies was not very homogeneous. The quality assessment identified some studies with low and average quality, which were excluded from the meta-analysis. We included only high-quality studies in our meta-analysis of RCT's and nRCT. The quality of the included observational studies is predominantly good. Table 2 A-D summarises the results of the quality appraisal of the included research papers.

### *Mask type*

Of the 37 meta-analytically evaluated studies, 31 examined the N95 mask, 19 the surgical mask with 1 not reporting on the specific type of mask due to the predominantly psychological research topic. There were 14 Studies evaluating both mask types (surgical and N95) and we compared the results in a separate meta-analysis (see below, Meta-analysis of N95 mask vs surgical mask).

### *Participants and time*

8641 subjects were used to conduct the meta-analysis totalling 22127 individual measurements/surveys.

This population consisted of young (age=34.8±12.5) and predominantly female subjects (m=2482, f=6159).

Physiological, physical and biochemical data was used in the meta-analyses comprising of 934 participants and 3765 experimental measurements.

The pooled prevalence data was drawn from a study population of n=8128 and included 17383 data entries.

Most of the 37 studies, evaluated in meta-analyses included healthy participants. Twelve studies were conducted in health care workers (32%).

Two studies (5%) included chronic obstructive pulmonary disease (COPD), one study on haemodialysis patients, another study included children (3%) and 4 studies involved pregnant women (11%).

The median experimental time of the studies included in the meta-analyses (mostly controlled trials) on physiological, physical, and chemical face mask effects was 18 minutes with an interquartile range (IQR) of 50 minutes (min.: 6 minutes, max.: 360 minutes). There was a major deviating mask exposure duration with exceptions (mean of 45.8 minutes with a standard deviation of 69.9 minutes). Therefore, the mean was not an appropriate parameter to characterise this distribution).

The study with the longest experimental duration (360 minutes, observational) included only 21 healthy participants, which corresponds to 2.2% of the total population studied (n=934).

Interestingly, the studies on symptoms (including many observational studies) had significantly longer observation times and a mean of 263.8±170.3 minutes (median 240, IQR 180) in a total of n=8128 participants.

**Table 2 A-D:** Summary of the quality appraisals for the included studies. Part A shows the quality analysis of RCTs with Cochrane RoB tool++, while Part B lists the results of the quality analysis of nRCTs with CASP checklist. Part C is on the quality analysis of observational (non questionnaire) studies with CASP checklist. Part D documents the quality analysis of the questionnaire studies by means of a similar checklist.

Table 2 A: Quality appraisal of randomised controlled trials

Publication	Selection Bias		Performance Bias	Detection Bias	Attrition Bias	Reporting Bias	7. Other Bias
	1. Random Sampling	2. Allocation Blinding	3. Blinding for Intervention	4. Evaluation Blinding	5. Incomplete Data	6. Selective Reporting	
Bertoli 2020	LR	LR	HR	HR	LR	UC	LR
Butz 2005	LR	LR	HR	LR	UC	UC	UC
Dirol 2021	LR	LR	HR	LR	LR	LR	LR
Fikenzer 2020	LR	LR	HR	LR	LR	LR	LR
Georgi 2020	LR	LR	HR	LR	LR	UC	LR
Goh 2019	LR	LR	HR	LR	LR	LR	LR
Hua 2020	LR	LR	HR	LR	UC	UC	LR
Kim J.H. 2013	HR	LR	HR	LR	LR	LR	LR
Kim J.H. 2015	LR	LR	HR	LR	LR	UC	LR
Kim J.H. 2016	LR	LR	HR	LR	LR	UC	LR
Mapelli 2021	LR	LR	HR	LR	LR	UC	LR
Roberge 2014	LR	LR	HR	LR	LR	UC	LR
Wong A.Y.-Y 2020	LR	LR	HR	LR	LR	UC	LR
Zhang 2021	LR	LR	HR	LR	LR	LR	LR
Legend: LR=low risk, HR=high risk, UC=Unclear							

Table 2 B: Quality appraisal of non-randomised controlled trials

<b>Publication</b>	<b>1. clear focus?</b>	<b>2. appropriate methods?</b>	<b>3. recruitment comprehensible?</b>	<b>4. valid measurement of exposure?</b>	<b>5. valid measurement of outcome?</b>	<b>6. equality of groups?</b>	<b>7. confounders taken into account?</b>	<b>8. sufficient size and significance of the effect?</b>	<b>9. credibility of the results?</b>	<b>10. transferability to other populations? clear focus?</b>	<b>11. comparability with existing evidence?</b>
Bharatendu 2020	Y	Y	Y	Y	UC	Y	Y	Y	UC	Y	UC
Coniam 2005	UC	N	Y	Y	Y	UC	UC	Y	Y	Y	UC
Epstein 2021	Y	Y	Y	Y	Y	Y	UC	N	Y	Y	Y
Lee 2011	Y	Y	Y	Y	N	Y	Y	N	UC	Y	UC
Roberge 2012	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	Y
Roberge 2010	Y	Y	Y	Y	Y	Y	Y	N	Y	Y	Y
Scarano 2020	Y	Y	Y	Y	Y	Y	UC	Y	Y	Y	UC
Shenal 2012	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	UC
Tong 2015	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	UC

Legend: Y=Yes, N=No, UC=Unclear

Table 2 C: Quality Appraisal of the Observational Studies

<b>Publication</b>	<b>1. clear focus?</b>	<b>2. appropriate methods?</b>	<b>3. recruitment comprehensible?</b>	<b>4. valid measurement of exposure?</b>	<b>5. valid measurement of outcome?</b>	<b>6. equality of groups?</b>	<b>7. confounders taken into account?</b>	<b>8. sufficient size and significance of the effect?</b>	<b>9. credibility of the results?</b>	<b>10. transferability to other populations? clear focus?</b>	<b>11. comparability with existing evidence?</b>
Beder 2008	Y	Y	N	Y	Y	UC	N	Y	Y	Y	Y
Choudhury 2020	Y	Y	Y	Y	Y	N	Y	Y	Y	N	N
Islam 2022	Y	Y	Y	Y	Y	Y	UC	Y	UC	Y	Y
Jafari 2021	Y	Y	Y	Y	Y	Y	UC	Y	Y	N	UC
Kao 2004	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	UC
Klimek 2020	Y	Y	Y	Y	Y	Y	UC	Y	Y	Y	UC
Kyung 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	UC
Li 2005	Y	Y	Y	Y	Y	Y	Y	UC	Y	Y	UC
Luckman 2020	Y	UC	N	Y	Y	Y	Y	Y	Y	Y	UC
Mo 2020	Y	Y	Y	Y	Y	UC	UC	Y	Y	Y	Y
Park 2020	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	UC
Pifarre 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	UC
Rebmann 2013	Y	Y	Y	Y	Y	Y	N	N	Y	N	Y
Sukul 2022	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	UC
Thomas 2011	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	UC
Toprak 2021	Y	Y	Y	Y	Y	UC	N	Y	Y	N	Y
Tornero-Aguilera 2021	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y

Legend: Y=Yes, N=No, UC=Unclear

Table 2 D: Quality Appraisal of the Questionnaire Studies

Publication	Study design	Validity and reliability	Questionnaire quality	Questionnaire design	Sample	Distribution and response	Analysis	Results	Summary and recommendation							
	Was a questionnaire study an appropriate method?	Are the results valid and realistic?	Does the questionnaire used provide reliable results?	Were sample questions provided?	Are the questions formulated in a clear and understandable way?	Details on how the questionnaire was prepared?	Was the questionnaire prepared in an appropriate manner?	Was the sample sufficiently large and representative?	Was information provided on how the questionnaire was made available?	Was information provided on response rates and exclusion criteria?	Was potential response bias discussed?	Were the results analysed appropriately?	Were all relevant results published?	Were both significant and non-significant results published?	Were results adequately interpreted?	Does the summary reflect the results of the study?
Foo 2006	Y	Y	UC	N	UC	UC	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
Forge 2009	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	UC	Y	N	Y	Y	Y
Heider 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Lan 2020	Y	Y	UC	N	UC	UC	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
Lim 2006	Y	Y	UC	N	UC	UC	Y	N	Y	N	Y	Y	Y	Y	Y	Y
Matusiak	Y	Y	UC	N	UC	Y	Y	Y	Y	N	Y	UC	Y	Y	Y	Y

*Consequences of wearing face masks*

2020																		
Naylor 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N
Ong 2020	Y	Y	UC	N	UC	N	UC	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	Y
Prousa 2020	Y	N	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	Y	Y	UC
Ramirez 2020	Y	Y	UC	N	UC	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	Y
Rosner 2020	Y	Y	UC	N	UC	N	UC	Y	Y	N	N	Y	Y	Y	Y	Y	Y	Y
Szczesniak 2020	Y	N	UC	N	UC	N	UC	Y	Y	N	N	Y	UC	N	N	Y	Y	Y
Szepietowski 2020	Y	Y	UC	N	UC	Y	Y	Y	Y	N	N	Y	Y	Y	Y	Y	Y	Y
Techasatian 2020	Y	Y	UC	N	UC	N	UC	Y	Y	N	N	Y	Y	Y	Y	Y	Y	Y
	Study design	Validity and reliability		Questionnaire quality		Questionnaire design		Sample	Distribution and response			Analysis	Results			Summary and recommendation		
Legend: Y=Yes, N=No, UC=Unclear																		

### Qualitative evaluation

Of the 54 included studies, 51 reported numerous adverse mask effects across multiple clinical disciplines, as already compiled in a previous scoping review<sup>8</sup>. Also fourteen of the 17 studies, which were not included in the meta-analysis reported those numerous mask effects.

Overall, our systematic review found mask related symptoms that can be classified under the previously described Mask-Induced Exhaustion Syndrome (MIES)<sup>8</sup>, with typical changes and symptoms that are often observed in combination.

Among the included 54 studies (Table 1), we detected and compiled reports on frequently statistically significant physiological and psychological changes ( $p < 0.05$ ) belonging to the MIES such as:

- increase in breathing dead space volume<sup>43,44</sup>
- increase in breathing resistance<sup>45-49</sup>
- increase in blood carbon dioxide<sup>20,43-45,50-64</sup>
- decrease in blood oxygen saturation<sup>20,44,45,48,49,52,53,55,56,58,60,61,63-68</sup>
- increase in heart rate<sup>20,44,47,49,52,56-58,61,62,65,67-69</sup>
- decrease in cardiopulmonary capacity<sup>45,48,64</sup>
- changes in respiratory rate<sup>44,45,48,52,53,57,58,62,64,66,67</sup>
- shortness of breath and difficulty breathing<sup>41,45,47,52,53,55,58,59,61,61,62,66,70-73</sup>
- headache<sup>50,53,61,70,72,74-78</sup>
- dizziness<sup>53,58,66</sup>
- feeling hot and clammy<sup>44,45,47,52,55,62,71,73</sup>
- decreased ability to concentrate<sup>68</sup>
- decreased ability to think<sup>58,61,68,78</sup>
- drowsiness<sup>78</sup>
- impaired skin barrier function<sup>41,78,79</sup>
- itching<sup>45,41,47,52,71,79-83,75</sup>
- acne, skin lesions and irritation<sup>79,41,58,71,62,78,82,75</sup>
- false sense of security<sup>84,85</sup>
- overall perceived fatigue and exhaustion<sup>70,52,45,53,66,55,56,47,48,44,62,86,63,69,64,61</sup>

Moreover, we could objectify additional symptoms of the MIES as follows:

- decrease in ventilation<sup>45,48,64</sup>
- increase in blood pressure<sup>20,45,47,48,52,58,59,64,67</sup>
- increase of measured temperature of the skin under the mask<sup>55,62,73,87</sup>
- increase of measured humidity of the air under the mask<sup>55,73,87</sup>
- communication disturbance<sup>61,71,78,88,89</sup>
- voice disorder<sup>71,90</sup>
- perceived discomfort<sup>41,45,52,73</sup>
- increased anxiety<sup>72,88,91</sup>
- increased mood swings or depressive mood<sup>72,88,90,91</sup>

and:

- changes in microbial metabolism<sup>20,92</sup>

However, three studies (6% of the included papers) describe the absence of adverse or even positive mask effects<sup>85,93,94</sup>.

## **Results of the meta-analysis**

In the meta-analytic evaluation, we found biochemical, physiological, physical, and perceptual symptoms with face mask use. We were also able to meta-analyse the pooled prevalence of symptoms. These results are presented in detail below.

### **Meta-analysis of biochemical effects of face masks**

#### *SpO<sub>2</sub> and face masks*

The results are summarised in Figure 2A.

In a pooled analysis, blood oxygen saturation is significantly lowered during mask use. This could be found for general mask use ( $p=0.0004$ ,  $SMD=-0.24$ , 95% CI -0.38 to -0.11,  $Z=3.53$ ,  $I^2=0\%$ ). The Eggers' test does not indicate the presence of funnel plot asymmetry ( $t(df=11)=-0.70$ ,  $p=.50$ ). This was also confirmed in the subgroup analysis for N95 mask use ( $p=0.001$ ,  $SMD=-0.3$ , 95% CI -0.49 to -0.12,  $Z=3.19$ ,  $I^2=0\%$ ), but not for surgical mask use ( $p=0.08$ ,  $SMD=-0.17$ , 95% CI [-0.37; 0.02],  $Z=1.77$ ,  $I^2=0\%$ ). However, 7 of 9 studies in the N95 mask meta-analysis contain the "0" in the confidence interval and are not significant because of  $n$  being too small (sample size). From the pooled analysis, it seems that N95 mask use may be responsible for a larger SpO<sub>2</sub> drop than surgical masks.

In a separate meta-analysis of pre-post studies an equally significant drop in SpO<sub>2</sub> was found when using a mask ( $p=0.0001$ ,  $SMD=-1.24$ , 95% CI -1.87 to -0.61,  $Z=3.87$ ,  $I^2=80\%$ ) especially in the subgroup of N95 masks ( $p=0.02$ ,  $SMD=-1.24$ , 95% CI -2.26 to -0.22,  $Z=2.37$ ,  $I^2=89\%$ ), yet with a high heterogeneity.

#### *Blood CO<sub>2</sub> content and face masks*

The results are summarised in Figure 2B.

In a pooled analysis, blood carbon dioxide content was found to be significantly elevated in mask use. This was perceived for general mask use ( $p=0.0001$ ,  $SMD=0.64$ , 95% CI 0.31 to 0.96,  $Z=3.86$ ,  $I^2=81\%$ ). The Eggers' test does not indicate the presence of funnel plot asymmetry ( $t(df=11)=-0.87$ ,  $p=.40$ ). This was also confirmed for N95 mask use ( $p=0.003$ ,  $SMD=0.78$ , 95% CI 0.28 to 1.29,  $Z=3.02$ ,  $I^2=84\%$ ) and also for surgical mask use ( $p<0.00001$ ,  $SMD=0.42$ , 95% CI 0.24 to 0.59,  $Z=4.65$ ,  $I^2=0\%$ ).

There was no significant difference between the pooled effect sizes of N95 and surgical masks ( $Q(df=1)=3.09$ ,  $p=.08$ ). Further separate pooled evaluations were also carried out for PtCO<sub>2</sub>, ETCO<sub>2</sub> and PaCO<sub>2</sub>, for each surgical and N95 masks with a significant increase in blood CO<sub>2</sub> with predominantly low heterogeneity.

Even in a separate meta-analysis of pre-post studies with high heterogeneity, a significant increase in blood carbon dioxide content was found when using a mask ( $p=0.003$ ,  $SMD=1.44$ , 95% CI 0.49 to 2.39,  $Z=2.97$ ,  $I^2=94\%$ ) and also in the subgroup of N95 masks ( $p=0.02$ ,  $SMD=1.51$ , 95% CI 0.24 to 2.78,  $Z=2.34$ ,  $I^2=96\%$ ).

Interestingly, 11 of 17 studies include "0" in the confidence interval and the majority showed no effect. The studies that showed clear effects (not including 0 in their confidence interval) differed from those that showed no certain effects as they either included N95 and/or pregnant women or children. The study by Dirol et al is an exception but has a sample size of  $n=100$  for surgical masks. Apparently, it takes N95 masks and vulnerable populations or appropriately large samples in surgical masks to make the effects more quantifiable and precise.

Accordingly, in the surgical mask meta-analysis, studies that included "0" in the confidence interval were of small sample size, with a mean of  $n=24$  and a median of  $n=14$ . The advantage of a meta-analysis is to combine several imprecise effects into a more precise overall effect<sup>30</sup>.

### **Meta-analysis of physiological effects of face masks**

#### *Ventilation( $V_E$ ) in L/min and face masks*

The results are summarised in Figure 3A.

Despite compensatory mechanisms, breathing volume (L/min) is significantly lowered during mask use in the pooled analysis.

This was not only verified for general mask use ( $p<0.00001$ , SMD= -0.72,  $Z=5.36$ , 95% CI -0.99 to -0.46,  $I^2=0\%$ ) in studies evaluated with an overall low heterogeneity ( $I^2=0$ ), but also for surgical ( $p<0.0001$ , SMD= -0.54, 95% CI -0.94 to -0.35,  $Z=4.32$ ,  $I^2=0\%$ ) and N95 mask use ( $p=0.0007$ , SMD= -1.06, 95% CI -1.68 to -0.45,  $Z=3.39$ ,  $I^2=0\%$ ). Both studies had an overall low heterogeneity( $I^2=0$ ).

On average, masks reduce respiratory minute volume by -19% according to our meta-analysis, and by as much as -24% for N95 masks; the difference between surgical and N95 masks is -10% respiratory minute volume.

#### *Respiratory rate and face masks*

The results are summarised in Figure 3B.

Interestingly, no statistical difference regarding respiratory rate was determined in mask use in the pooled analysis.

Even in the subgroups containing N95 and surgical masks, no difference compared to the no mask condition could be found.

#### *Systolic blood pressure (SBP) and masks*

The results are summarised in Figure 4A.

A significant elevation in systolic blood pressure was found for mask users with  $p=0.02$ , SMD= 0.17, 95% CI 0.03 to 0.32,  $Z=2.39$  and  $I^2=0\%$  in the pooled analysis. It is a small effect and in 9 out of 10 studies insignificant, including 2 with higher  $n$  in each case. The Eggers' test does not indicate the presence of funnel plot asymmetry ( $t(df=8)$ ,  $p=.27$ ). This was verified in the subgroup analysis for surgical masks ( $p=0.02$ , SMD= 0.21, 95% CI 0.03 to 0.39,  $Z=2.33$ ,  $I^2=0\%$ ). In studies evaluating both mask types (surgical and N95) the N95 mask always yielded a higher SBP than the surgical mask. However, this effect was not statistically significant. There is no significant difference between the pooled effect sizes of N95 and surgical masks ( $Q(df=1)=0.98$ ,  $p=.32$ ).

#### *Heart rate and masks*

The results are summarised in Figure 4B.

No statistically significant difference regarding the heart rate during mask use was found in the pooled analysis. The Eggers' test does not indicate the presence of funnel plot asymmetry ( $t(df=14)$ ,  $p=.94$ ). However, in the subgroup analysis containing surgical and N95 masks, only for the N95 mask condition a weak significance for a slight increase in heart rate could be found ( $p=0.02$ , SMD= 0.22, 95%CI 0.03 to 0.41,  $Z=2.30$  and low heterogeneity of studies with  $I^2=0$ ). There is no significant difference between the pooled effect sizes of N95 and surgical masks ( $Q(df=1)=1.26$ ,  $p=.26$ ).

### ***Meta-analysis of physical effects of face masks***

#### *Skin temperature and face masks*

The results are summarised in Figure 5A.

Skin covered by mask has a significantly higher temperature during rest and activity. This could be found for general mask use ( $p=0.005$ ,  $SMD=0.80$ , 95% CI 0.23 to 1.38,  $Z=2.81$ ,  $I^2=72\%$ ), for N95 mask use ( $p=0.02$ ,  $SMD=0.72$ , 95% CI 0.12 to 1.32,  $Z=2.35$ ,  $I^2=55\%$ ), but not for surgical mask use ( $p=0.21$ ,  $SMD=0.96$ ,  $Z=1.26$ ,  $I^2=90\%$ ).

#### *Humidity and face masks*

The results are summarised in Figure 5B.

The dead space covered by mask has a significantly higher humidity in the pooled analysis. This could be found for general mask use with  $p<0.00001$ ,  $SMD=2.24$ , 95% CI 1.32 to 3.17,  $Z=4.75$  and  $I^2=50\%$ ).

### ***Meta-analysis of measured symptoms and sensations during face mask use***

#### *Discomfort and face masks*

The results are summarised in Figure 6A.

Perceived discomfort is significantly higher in mask use during rest and activity in the pooled analysis.

This could be found for general mask use ( $p<0.0001$ ,  $SMD=1.16$ , 95% CI 0.58 to 1.73,  $Z=3.94$ ,  $I^2=74\%$ ), for N95 mask use ( $p<0.00001$ ,  $SMD=1.98$ , 95% CI 1.37 to 2.59,  $Z=6.34$ ,  $I^2=0\%$ ) as well as for surgical mask use ( $p<0.00001$ ,  $SMD=0.71$ , 95% CI 0.46 to 0.96,  $Z=5.58$ ,  $I^2=0\%$ ).

#### *Itch and face masks*

The results are summarised in Figure 6B.

In N95 mask use, the perceived itching was significantly elevated ( $p=0.003$ ,  $SMD=2.65$ , 95% CI 1.21 to 4.09,  $Z=3.6$ ,  $I^2=83\%$ ) during activity according to the pooled subgroup analysis.

Although not statistically significant, an overall tendency for itching was found for general mask use in the pooled analysis.

#### *Exertion and face masks*

The results are summarised in Figure 6C.

Perceived exertion is significantly higher in mask use during activity in the pooled analysis.

This could be found for general mask use ( $p<0.0001$ ,  $SMD=0.90$ , 95% CI 0.58 to 1.23,  $Z=5.31$ ,  $I^2=71\%$ ), for N95 mask use ( $p=0.002$ ,  $SMD=1.19$ , 95% CI 0.43 to 1.95,  $Z=3.06$ ,  $I^2=81\%$ ) as well as for surgical mask use ( $p<0.0001$ ,  $SMD=0.63$ , 95% CI 0.40 to 0.87,  $Z=5.29$ ,  $I^2=24\%$ ). The Eggers' test indicates the presence of funnel plot asymmetry ( $t(df=10)=2.68$ ,  $p=.02$ ). For N95 mask use ( $p=0.002$ ,  $SMD=1.19$ ,  $Z=3.06$ ,  $I^2=81\%$ ) and this result was confirmed for surgical mask use too ( $p<0.0001$ ,  $SMD=0.63$ ,  $Z=5.29$ ,  $I^2=24\%$ ). There is no significant difference between the pooled effect sizes of N95 and surgical masks ( $Q(df=1)=1.97$ ,  $p=.16$ ).

#### *Shortness of breath and face masks*

The results are summarised in Figure 6D.

Perceived shortness of breath is significantly higher during mask use during activity in the pooled analysis ( $p=0.006$ ,  $SMD=1.46$ , 95% CI 0.42 to 2.50,  $Z=2.75$ ,  $I^2=86\%$ ).

In the subgroup analysis for surgical and N95 masks, the masks always led to an increase in perceived shortness of breath, but the number of studies that could be included was very limited and no statistically significant results were found in the subgroup analysis.

#### *Perceived heat and face masks*

The results are summarised in Figure 6E.

Perceived heat is significantly higher during mask use with physical activity in the pooled analysis ( $p=0.002$ ,  $SMD=0.70$ , 95%CI 0.28 to 1.13,  $Z=3.27$ ,  $I^2=62\%$ ).

In the subgroup analysis containing surgical and N95 masks the heat perception was increased in both mask types, but only for the surgical mask condition a statistical significance for an increase in heat perception could be found ( $p=0.008$ ,  $SDM=0.61$ , 95% CI 0.16 to 1.06,  $Z=2.66$ ,  $I^2=50\%$ ).

#### *Perceived humidity and face masks*

The results are summarised in Figure 6F.

Perceived humidity is significantly higher in mask use during activity according to the pooled analysis ( $p=0.002$ ,  $SMD=0.90$ , 95% CI 0.34 to 1.46,  $Z=3.17$ ,  $I^2=53\%$ ).

The subgroup analysis containing surgical and N95 masks was completed merely for surgical masks due to lack of studies on N95 masks.

In the surgical mask condition a statistical significance for an increase in humidity perception could be found ( $p<0.00001$ ,  $SMD=0.63$ , 95% CI 0.36 to 0.90,  $Z=4.6$ ,  $I^2=0$ ).

#### ***Meta-analysis of N95 mask vs surgical mask***

The results are summarised in Figure 7A-C.

The N95 mask leads to measurably worse effects compared to the surgical mask. The blood oxygenation is significantly decreased when using a N95 mask compared to a surgical mask with  $p=0.003$ ,  $SMD= -0.53$ , 95%CI -0.88 to -0.18,  $Z=2.98$ ,  $I^2=37\%$ . The heart rate ( $p=0.01$ ,  $SMD=0.25$ , 95% CI 0.05 to 0.45,  $Z=2.47$ ,  $I^2=0\%$ ), the perception of discomfort ( $p=0.02$ ,  $SMD=3.07$ , 95% CI 0.52 to 5.61,  $Z=2.36$ ,  $I^2=95\%$ ) and humidity ( $p=0.02$ ,  $SMD=0.59$ , 95% CI 0.09 to 1.10,  $Z=2.32$ ,  $I^2=0\%$ ) increased in each case when the N95 mask was compared to the surgical mask. This trend was also evident for  $CO_2$ , minute volume, exertion, heat, shortened breath and systolic blood pressure, but was not statistically significant due to the limited available studies.

#### ***Meta-analysis with pooled prevalence of symptoms during face mask use***

The results are summarised in Figure 8.

The prevalence of headaches with mask use is significant in the majority of evaluated users ( $n=2525$ ), with a pooled prevalence of 62% ( $p<0.00001$ , 95%CI 0.48 to 0.75) and even 70% ( $p<0.00001$ , 95%CI 0.52 to 0.88) with N95 mask use.

Acne when using a mask is significantly present in evaluated users ( $n=1489$ ) with a pooled prevalence of 38% in general mask use ( $p<0.00001$ , 95%CI 0.22 to 0.54).

Skin irritation occurrence when using a mask is significantly present in the evaluated users ( $n=3046$ ) with a pooled prevalence of 36% in general mask use ( $p<0.00001$ , 95%CI 0.24 to 0.49).

Shortness of breath rate when using a mask is significantly present in users (n=2134) with a pooled prevalence of 33% in general mask use ( $p < 0.00001$ , 95%CI 0.23 to 0.44) and 37% in N95 mask use ( $p = 0.01$ , 95%CI 0.07 to 0.67).

The prevalence of itch with mask use is substantial in evaluated users (n=5000), with a pooled prevalence of 26% ( $p < 0.00001$ , 95%CI 0.15 to 0.36). In the subgroup analysis, the pooled prevalence for itch in N95 mask use was 51% ( $p < 0.00001$ , 95%CI 0.47 to 0.55) while it was 17% in surgical mask use ( $p < 0.0001$ , 95%CI 0.09 to 0.26). These results were confirmed in control calculations using the R software.

The prevalence of voice disorders when using a mask is significant in evaluated users (n=1097) with a pooled prevalence of 23% in general mask use ( $p = 0.03$ , 95%CI 0.02 to 0.43), however with high heterogeneity of the included studies.

The prevalence of dizziness when using a mask is significant in evaluated users (n=153) with a pooled prevalence of 5% in general mask use ( $p = 0.01$ , 95%CI 0.01 to 0.09). Due to the small sample size (n) in the referred studies, there are wide confidence intervals. This results in a significant, but not really pronounced overall result for dizziness.

## Discussion

Besides the anticipated protection against the transmission of pathogens, face masks undoubtedly impede natural breathing. Such respiratory impairments due to the “new-normal” lifestyle under the present global pandemic have imposed potential adverse effects on our usual external and internal respiration, affecting a wide range of physio-metabolic processes within various organ systems and/or at cellular levels<sup>8,20</sup>. Ensuing consequences were eventually observed at the physical, psychological and social levels along with certain clinical symptoms in the individual human being<sup>8</sup>. In this systemic review, we applied meta-analysis and comprehensive evaluations of physio-metabolic, physical, psychological and clinical burdens of wearing face masks in the general population. Restricting breathing through face masks has turned out to be a fundamental, incisive intervention with possible negative effects on public health.

### *Physio-metabolic burden of masks*

Our meta-analysis clearly depicts that masks significantly restrict O<sub>2</sub> uptake and hinder CO<sub>2</sub> release. Based on the meta-analytic effect sizes defined by Cohen<sup>95</sup>, the effect size for CO<sub>2</sub> retention (as per PtCO<sub>2</sub>, ETCO<sub>2</sub> and PaCO<sub>2</sub> outcomes) is medium for all mask types and is larger for N95 masks. The effect size for O<sub>2</sub> uptake disturbance (as per SpO<sub>2</sub> outcome) is relatively smaller but highly significant ( $p = 0.0004$ ) (Fig. 9A and Fig. 2 A, B). Such respiratory gas-exchange discrepancy can be attributed to the constantly increased dead space ventilation volume<sup>8,43,44,96,97</sup> (i.e., continuous rebreathing from the masks dead space volume) and breathing resistance<sup>8,45-49</sup>. Continuous CO<sub>2</sub> rebreathing causes the right-shift of haemoglobin-O<sub>2</sub> saturation curve. Since O<sub>2</sub> and CO<sub>2</sub> homeostasis influences diverse down-stream metabolic processes, corresponding changes towards clinically concerning directions may lead to unfavourable consequences such as transient hypoxaemia and hypercarbia, increased breath humidity and body temperature along with compromised physiological compensations etc..

### *Transient hypoxaemia*

A progressive decrease in SpO<sub>2</sub> is observed with respect to the duration of wearing a mask<sup>20,52,55,57,58,60,65,70,98</sup>. The decline in SpO<sub>2</sub> levels confirmed in our systemic-review supports the onset and progression of oxidative stress (via significantly increased exhaled breath aldehydes – originating from lipid peroxidation) reported by Sukul et al<sup>20</sup>. Studies have shown that oxidative

stress (under hypoxic conditions) can inhibit cell-mediated immune response (e.g. T-lymphocytes, TCR CD4 complex etc.) to fight viral infections, which may gradually lead to immune suppression<sup>99,100</sup>. Arterial hypoxaemia increases the level of the hypoxia inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ), which further inhibits T-cells and stimulates regulatory T-cells<sup>100</sup>. This may set the stage for contracting any infection, including SARS-CoV-2 and making the consequences of that infection much more severe. In essence, masks may put wearers at an increased risk of infection and severity<sup>99–101</sup>. A recent review<sup>102</sup> by Serebrovska et al discusses a possible link between HIF-1 $\alpha$  activation and cell entry of SARS-CoV-2. If the cell is already under oxidative stress, activation of HIF-1 $\alpha$  may suppress important adaptive mechanisms e.g., autophagy or proteasomal proteolysis is leads to the induction of necrosis and excessive cytokine production. Sturrock et al<sup>103</sup> demonstrated that the SARS-CoV-2 receptor (e.g., ACE2 and TMPRSS2) expression by primary type II alveolar epithelial cells increased significantly following exposure to hypoxic environments *in vivo* and *in vitro*. Furthermore, recent research has demonstrated that the cellular entry of SARS-CoV-2 also depends on many other receptor paths/routes (e.g., CD147, CD147 - spike proteins etc.), mediated by HIF-1 $\alpha$  upregulation<sup>104–107</sup>. Therefore, the effect of even mild hypoxaemia for an extended span may promote an infection risk along with metabolic stress e.g., due to altered pH via respiratory acidosis. In line with that, Sukul et al<sup>20</sup> observed a significant decrease in exhaled volatile metabolites (e.g. organosulfur and short-chain fatty acids) originating from the lower gut microbiota during face mask use – indicating anaerobiosis, metabolic acidosis and possible immunosuppression. Even marginal local effects of masks on salivary metabolites in young and healthy adults have indicated alteration of microbial metabolic activity<sup>92</sup>.

The findings of Spira 2022<sup>10</sup> from European data show that mask use correlates with increased morbidity and mortality, which could be due to the above-discussed possible processes. Moreover, prolonged hypoxic conditions and low oxygen levels pave the way for immunosuppression and inflammation, which can promote the growth, invasion and spread of cancers<sup>107–109</sup>.

However, further experimental studies are needed to prove that hypoxaemia under long-term mask use may result in quantifiable changes in HIF-1 $\alpha$  and immunosuppression – especially in elderly, ill/comorbid and/or immunocompromised individuals.

### *Transient hypercarbia*

In line with the increased dead space ventilation and consistently decreasing SpO<sub>2</sub> level, CO<sub>2</sub> inhalation elevates progressively during the course of wearing a mask, causing transient hypercarbia<sup>20,52,55,57,58,60,98</sup>. Very recent experimental data exist on CO<sub>2</sub> concentrations of concern in the air breathed while wearing masks, especially in children<sup>110,111</sup>. Systemic CO<sub>2</sub> concentration exerts an important influence on the intra- and extracellular pH. CO<sub>2</sub> passes quickly through the cell membranes to form carbonic acid, which releases protons and in excess causes acidosis<sup>112–114</sup>. With a prolonged CO<sub>2</sub> burden the body uses the bones (CO<sub>2</sub> storage) to regulate the blood pH: bicarbonate and a positive ion (Ca<sup>2+</sup>, K<sup>+</sup>, Na<sup>+</sup>) are exchanged for H<sup>+</sup>. Accordingly, kidney and organ calcification were frequently seen in animal studies on low-level CO<sub>2</sub> exposure<sup>115,116</sup>. Additionally, CO<sub>2</sub> in relationship with chronic and/or intermittent long-term exposure might induce pathological states by favouring DNA alterations and inflammation<sup>117,118</sup>. Moreover, inflammation is reported to be caused by low-level CO<sub>2</sub> exposure in humans and animals<sup>118–122</sup>. Even slightly elevated CO<sub>2</sub> induces higher levels of pro-inflammatory Interleukin-1 $\beta$ , a protein involved in regulating immune responses, which causes inflammation, vasoconstriction and vascular damage<sup>121</sup>. In addition, carbon dioxide is also known as a trigger of oxidative stress caused by reactive oxygen species (ROS)<sup>117</sup> including oxidative damage to cellular DNA<sup>117,118</sup>.

Altogether, the possible damaging mechanism of CO<sub>2</sub> affecting tissues is based on the conditions of oxidative stress and acidosis with increased inflammation and apoptosis as described above<sup>117,119–124</sup>. In the long term, therefore, this could be possible during mask use even at blood-CO<sub>2</sub> levels that do

not reach the thresholds. In spontaneously breathing subjects in a sitting position, exhaled CO<sub>2</sub> profiles mirror the endogenous isoprene exhalation<sup>12,125</sup>. Significant and progressively decreased breath isoprene recently observed in adults<sup>20</sup> indicates the deoxygenation driven sympathetic vasoconstriction in the peripheral compartments<sup>126</sup>. Prolonged deoxygenation and CO<sub>2</sub> re-breathing therefore, may eventually lead to pulmonary vasoconstriction that may hinder blood-CO<sub>2</sub> levels to reach the thresholds. For instance, Sukul et. al also reported the presence of significant hyperventilation state in older adults aged  $\geq 60$  years before wearing a face mask for the participation in experiments. This indicates a compromised respiratory compensation of precedent mask use (which was obligatory due to pandemic regulations at that time) by these subjects.

#### *Physical burden of masks: Humidity and skin temperature*

Together with the immune-inhibiting mechanisms mentioned above, we found some other possible deleterious mask effects that contradict healthy natural breathing. The most prominent and extreme effect was found in the increase of air humidity and skin temperature within the dead space of the mask (Figure 9B and Figure 5). Increased humidity and temperature can increase droplet and aerosol generation, which facilitate liquid penetration through the mask mesh. This not only increases the chance of microorganism (fungal and bacterial pathogens) growth on and in masks<sup>127-129</sup> causing increased risk for accumulation of fungal and bacterial pathogens<sup>127,129</sup> including mucormycosis<sup>130</sup>, but also leading to re-breathing of viruses that may be trapped and enriched within the moisturised mask meshwork. Therefore, these conditions within masks are favourable for pathogenic growth and are unfavourable for good/systemic microbiota i.e., individual specific. As a result, the isolation of people with masks for extended periods can attain conditions for new and individual specific strains formations/mutations of pathogens – to which other people in the environment will be susceptible and/or not immune. Additionally, the high concentration of microbiome in masks can be a potential source of infection for the population. The findings of Fögen 2022<sup>5</sup> using data from the USA which shows that mask use correlates with an increased mortality could be due to these processes. This phenomenon could also explain the similar figures found by Spira<sup>10</sup> in the EU.

#### *Compensatory physiological mechanisms*

Our meta-analytically quantified CO<sub>2</sub>-rise and O<sub>2</sub>-depletion (Figure 2, 9A) with mask use certainly needs physiological compensations (Figures 3, 4 and 10). Interestingly, the compensatory responses to mask wearing (e.g., rise in heart rate, changes in respiratory rate and/or minute ventilation etc.) was lower (absent or even reverse) than expected<sup>115,131,132</sup>. In former human experiments with low level 1-2% CO<sub>2</sub> exposure to breathing air – which corresponds to measured values during mask use<sup>133</sup> – an increased respiratory minute volume (V<sub>E</sub>) of >34% was detected<sup>115</sup>. In contrast to that and according to our results under masks a significantly decreased V<sub>E</sub> by -19% on an average and up to -24% under N95 masks occurs despite face mask driven CO<sub>2</sub> exposure<sup>133</sup>. Even the V<sub>E</sub> differed by 10% between N95 and surgical masks (Figure 3A). However, it appears to have no acute clinical impact in the short term and does not exceed normal values of SpO<sub>2</sub> and systemic CO<sub>2</sub> although these may become problematic in the long run. A compensatory higher arterial PaCO<sub>2</sub> and bicarbonate levels execute the buffering of inhaled CO<sub>2</sub>. Interestingly, during chronic breathing of low CO<sub>2</sub> concentrations (in the no-mask condition), due to compensatory mechanisms, e.g. lowered blood pH, increased respiratory rate and V<sub>E</sub><sup>115</sup> and an acclimatisation occurs<sup>115,131,132,134,135</sup>. In mask users, those compensatory mechanisms however seem to differ or get disturbed (e.g. no rise in respiratory rate, heart rate and simultaneous fall in V<sub>E</sub>). Health risks should be considered despite the mask related compensation attempts<sup>133</sup>. During face mask use a rise in the arterial PaCO<sub>2</sub> is possible in the long term<sup>20,52,58,60,98</sup>. Although, PaCO<sub>2</sub> generally remains at a sub-threshold level in

healthy mask users<sup>98,131</sup>, concerning pathological changes can occur in older (>60 years) and sick people<sup>20,59</sup>.

Our findings depicted an absence of typical compensatory reactions to transient hypercarbia thereby implying a suppression of a physiological response owing to the unusual conditions of wearing a mask. The reasons behind this phenomenon, i.e. the absence of a rise in the respiratory rate and ventilation, remain unclear. The simultaneous change in the adverse direction (CO<sub>2</sub> rise and simultaneous O<sub>2</sub> fall with concomitant dead space- and resistance enlargement caused by the mask) may be responsible for this. The drop in SpO<sub>2</sub> and the rise in CO<sub>2</sub> (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>) with no major changes in the heart rate in our meta-analysis also transpires to be an unexpected reaction. Sukul et al<sup>20</sup> reported altered breathing patterns, respiratory resistance and discomfort under medical masks. Adults younger than 60 years of age described slow breathing (slow and deep inspiration and expiration) under masks, whereas shallow/thoracic breathing (breathing with increased inhalation duration and effort), respiratory resistance and dyspnoea was portrayed by those ≥ 60 years of age. Fittingly, altered breathing patterns/kinetics, progressive changes towards deoxygenation, hypercarbia and insignificant changes in the respiratory and heart rate transpired to be surprising mask outcomes in our present results (hypercapnia-like effects). Thus, prolonged masks use may lead to hypercapnic hypoxia like conditions. While short and acute hypercapnic hypoxia like conditions in healthy individuals can promote positive effects (sport, training etc.), a chronic/prolonged hypercapnic hypoxia (as known from sleep apnoea) is toxic for the cardiovascular system in the long run – causing metabolic syndrome<sup>8</sup> as well as additional effects on cognitive functions<sup>136</sup>.

#### *N95 mask compared to surgical mask*

In line with recent findings by Kisielinski 2021<sup>8</sup> and Sukul 2022<sup>20</sup>, the present results clearly show that N95 masks lead to significantly more pronounced and unfavourable biochemical, physiological and psychological effects (Figure 7) than surgical masks. Altogether, the results in blood oxygenation, discomfort, heart rate, CO<sub>2</sub>, exertion, humidity, blood pressure, V<sub>E</sub>, temperature, dyspnoea and itching etc. can be attributed to the larger (almost doubled) dead space and higher breathing resistance of the N95 mask<sup>8</sup>. Compared to the surgical mask upon the short-term effects, N95 masks could impose elevated health risks under extended use. Interestingly, recent data from a large multi-country RCT study show no significant differences between the two mask types in terms of SARS-CoV2 infection rates<sup>137</sup>.

#### *Short mask experiment times*

It is noteworthy to say that in studies with short assessment times neither correspond to real-life conditions nor do they exclude short-term compensatory mechanisms, e.g. obvious for CO<sub>2</sub>-rebreathing. However, immediate compensatory mechanisms can hide further adverse reactions<sup>115,131,133</sup>. Therefore, longer observation times can lead to clearer values that are closer or above the thresholds due to the attenuation or collapse of transient physiological mechanisms. The experimental studies used here examined important outcomes only had a median examination time of 18 minutes (Figure 11). Heterogeneous studies with small sample sizes yielded significant and medium to strong results (Figures 10 and 12). Nevertheless, experimental studies with longer assessment periods are needed.

The observational studies included in the present analysis on symptoms were conducted over significantly longer periods (median 240 min, IQR 180) and are able to consider cumulative and long-term effects. It is known that observational studies are far more precise in finding negative effects and are particularly suitable to investigate exposures (e.g., air pollution or smoking) that are difficult or impossible to investigate in randomised controlled trials (RCTs). In addition,

observational studies are important to investigate causes with a long latency period, such as toxicological and carcinogenic effects from environmental exposures or drugs <sup>42</sup>. The longest period of included studies was 8 months with an averaged of wearing the mask 8 hours per day (observational study), however with the shortest study with a 5 minutes examining/exposition time (controlled trial).

*Possible sub-threshold impact of masks –the low-dose long-term effect on health*

In contrast to our study, most of the recent systematic reviews <sup>21–25</sup> have only analysed a few outcome threshold values without considering comprehensive effects, exposure time and the susceptibility of the exposed organisms and tissues. Therefore, their recommendations e.g. masks are harmless and safe for everybody etc. appears to be superficial, non-medical, non-holistic and misleading.

In accordance with conclusions of Fikenzler, Sukul and Zhang <sup>20,45,64</sup>, we have found hints to deleterious effects even without exceeding physiological threshold values and we have interpreted these data as a risk for individuals with suppressed compensatory mechanisms such as in elderly and sick subjects with cardiorespiratory diseases, infection, diabetes, cancer and other comorbidities. Sukul et al <sup>20</sup> were able to show that the unfavourable effects are more pronounced in the elderly (aged: 60 – 80 years). Moreover, they could provide evidence for toxic effects of face masks including oxidative stress, immunosuppression, deoxygenation and hypercarbia induced vasoconstriction and altered systemic microbial activity. Even with CO<sub>2</sub> and SpO<sub>2</sub> levels that do not exceed the limits, many clinical researchers have also found troubling results in face mask wearers. Neurologists observed changes in MRI brain signal baseline level due to face mask use <sup>9</sup>. Wearing a surgical mask for merely 9 minutes increased end-tidal CO<sub>2</sub> causing mild hypercapnia. This was responsible for a compensatory increase in cerebral blood flow with morphological changes similar to that of a CO<sub>2</sub> gas challenge or holding your breath. In patients with aneurysms or brain tumours this phenomenon could be deleterious. Another study showed a pathologic and altered brain metabolism while wearing a N95 mask for 6 hours <sup>11</sup>. The MRI imaging revealed a significant drop in brain oxygenation. A more than 50% drop in oxygenation in the cingulate gyrus (cognition circuit) after 6 hours of mask use was associated with clinical symptoms of a confused state in 80% of the subjects above 35 years. The authors even concluded that the general population should not wear a N95 mask. This phenomenon of brain deoxygenation could be dangerous for people with altered brain functions when on medication, after a transient ischaemic attack (TIA) or stroke respectively.

Ophthalmological studies indicated risk of retinal damage from long-term use of masks. N95 masks reduced the vascular density in the vascular plexus even under resting conditions as early as after 60 minutes <sup>138</sup>. Here, the drop in SpO<sub>2</sub> and increase in blood pressure were significant but within the normal physiological range. Another study reported a significant mask-induced increase in intraocular pressure (IOP) after approx. 5 minutes of wearing <sup>6</sup>. Thus, wearing masks may counteract the therapy aiming to reduce the IOP and can exacerbate irreversible long-term vision problems in individuals with glaucoma. Numerous other studies have shown that the long-term effects, leading to deleterious clinical outcome may result from prolonged mask wearing <sup>9,11,138,139</sup>. Such effects are comparable to sick building syndrome (SBS) <sup>140</sup>, cigarette smoking and other chronic, slightly toxic influences relevant to the general population.

In accordance with our present analysis and precedent scoping review <sup>8</sup>, mask-related changes in leaning towards pathological values can lead to illness and clinical consequences, just like chronically, repeated subliminal harmful environmental events. Occupational diseases defined by the International Labour organization (ILO) and that are in accordance with the worker's compensation act in Germany illustrates the potential harm caused by chronic exposure to subthreshold environmental factors <sup>141</sup>. Numerous examples of these principles can be found in the literature concerning pharmacology, toxicology, clinical and occupational medicine and even in

psychology<sup>142–151</sup>. Many other toxicological and environmental health examples are presented in the recent scoping review by Kisielinski et al<sup>8</sup>, which refers to MIES (Mask-Induced Exhaustion Syndrome). Such subliminal chronic changes and harmful effects in the long run are comparable to the sick building syndrome (SBS)<sup>140</sup>, cigarette smoking<sup>152</sup>, salty diet<sup>153</sup>, aluminium environmental pollution<sup>154</sup>, low-level lead exposure<sup>155</sup>, organochlorine pesticides and polychlorinated biphenyl exposure<sup>156</sup> or even the so-called climate change exposure<sup>157</sup>. Altogether, even the subliminal changes due to face mask use can become clinically relevant.

#### *Overlapping of face mask effects (MIES) with long-COVID symptoms*

Regarding the numerous mask symptoms an important question arises: Can masks be responsible for a misinterpreted long-COVID-syndrome after an effectively treated COVID-19 infection? Nearly 40% of main long-COVID symptoms<sup>158</sup> overlap with mask related complaints and symptoms described by Kisielinski et al as MIES<sup>8</sup> like fatigue, dyspnoea, confusion, anxiety, depression, tachycardia, dizziness, headache, which we also detected in the qualitative and quantitative analysis of face mask effects in our systematic review. It is possible that some symptoms attributed to long-COVID are predominantly mask-related. Further research on this phenomenon needs to be conducted.

#### *Complaints and symptoms under mask use and the WHO definition of health*

Amongst the perceived sensations with mask use only 6 symptoms (exertion, discomfort, shortness of breath, humidity, heat and itch) could be meta-analysed and have resulted in predominantly strong effect sizes (Figure 12). In the pooled prevalence analysis, we included eight main symptoms namely headache, acne, skin irritation, shortness of breath, heat, itch, voice disorder and dizziness (Figure 13) out of which all were significant in the evaluated population (Figure 8). There are many more reported in the literature. However, these could not be meta-analysed due to the low number of comparable studies on those particular complaints. In the included literature reported mask related symptoms were: rhinitis<sup>80</sup>, disabilities to think and to concentrate<sup>58,61,68,78</sup>, drowsiness<sup>78</sup>, communication disorder<sup>61,88,89</sup>, depression and mood swings<sup>72,88,90,91</sup>, anger<sup>72</sup>, perceived discomfort<sup>41,45,52,73</sup>, anxiety<sup>72,88,91</sup>, and an overall perceived fatigue and exhaustion<sup>44,45,47,48,52,53,55,56,61–64,66,69,70,86</sup>.

All of these mask-related symptoms contradict a state of well-being and health as defined by the WHO. According to the WHO; “health is a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity”<sup>159</sup>. Based on the facts we have found, the use of face mask in the hope of maintaining health is unfortunately contradicting the WHO's definition of health. Regarding all the possible side effects of mask and their still unproven efficacy against viral transmission within the general population<sup>4,160–162</sup>, health seems not to be substantially preserved by wearing face masks. So far, only two randomized controlled mask trials for prevention of SARS-CoV-2 infection in the general population have been published: one high quality study from Denmark, Europe<sup>163</sup>, and the other from Bangladesh with biased results and a lot of inconsistencies<sup>164</sup>. Based on a Bayesian random-effects meta-analysis of these two trials, the posterior median for relative risk was 0.91 (95% credible interval 0.63 to 1.33, 73% probability of some benefits with very limited evidence)<sup>165</sup>. The paucity in high-quality mask studies is unfortunate. Seeing the overall weak evidence for efficacy of masks against viral transmission within the general population<sup>4,160–162,166–168</sup>, face masks have to be evaluated appropriately in the sense of the *Hippocratic Oath* and as per the *Primum nihil nocere* (above all do not harm). To avoid at all costs that the damage caused by preventive or therapeutic measures becomes greater than that caused by the disease itself, should be the credo of all those involved in the containment of the crisis, including politicians and the so-called experts. Medical decisions can only be made on the basis of comprehensive knowledge on a patient's overall condition, individualised case history, considering all previous illnesses and

interventions, physical and mental predispositions and his/her socio-economic state etc. When it comes to medical decision-making in a sick person, the weighing of therapeutic measures for the benefit of the patient against the side effects of the therapy is to be evaluated differently than a prophylactic procedure in healthy people. If wrong decisions are made in the selection of preventive measures in healthy individuals, or if they are improperly applied, the consequences are usually much more severe and liability claims are often unavoidable. From a standardisation point of view the filtration efficacy of mask for viruses remains hypothetical and not in line with the established standards. There are national and international standards for bacteria filtration efficiency (BFE) for medical masks since decades, for example the EU-EN 14683, or the USA-ASTM F2101. They are the prerequisites for general approval. However, since 2020 (i.e. nearly 3 years), no comparable standard/testing of masks for viruses does yet exist. Given the fact, that medical masks (surgical and N95) increase particle exhalation in the smallest size range of 0.3 – 0.5  $\mu\text{m}$ , shifting the geometric mean diameter toward smaller sizes (longer in air) compared to no mask conditions<sup>169</sup> doubts arise. Such scientific facts are pointing towards the nebulisation effect of masks, which could be an add-on for their weakness against viral transmission in general.

### **Limitations**

While looking at the potential limitations our systematic review rarely discussed the inhaled toxins associated with the mask. Inhalation and ingestion of toxic substances, which are ingredients of the masks, are also of importance in evaluating this pandemic non-pharmaceutical intervention (NPI). In addition, our work has not extensively studied the microbial colonisation of masks and the consequences of contamination by microorganisms for the wearer. Based on the studies conducted during the pandemic, the control groups without masks were mostly the same individuals, or individuals who were not mask abstinent for too long (general mask requirement)<sup>170</sup>, so the mask-no-mask differences may be mitigated. Because of the rapid flow of science, new interesting papers have certainly appeared that we were unable to consider in the meta-analysis as they appeared after the period of our data search (search limitation to 31.12.2021). The most important and relevant observational studies were considered for this analysis thereby addressing the physio-metabolic and clinical effects. Numerous psychological and other effects could not be assessed analytically as too few relevant and evaluable studies were available.

### **Conclusion**

This systematic review comprehensively revealed ample evidence for multiple adverse physio-metabolic and clinical outcomes of medical face masks. This can have long-term clinical consequences, especially for vulnerable groups e.g. children, pregnant, elderly and ill. The N95 masks lead to measurably more adverse results than surgical masks. Besides transient and progressive hypoxaemia, hypercarbia and individualised clinical symptoms our findings are in line with reports on face masks driven down-stream aberrations (e.g. oxidative stress, hypercapnia, vasoconstriction, pro-inflammatory response, immunosuppression etc.) at the organ, cellular and microbiome levels and support the MIES (Mask Induced Exhaustion Syndrome). From our point of view, while a short application of the mask seems to be less harmful, longer and long-term use may cause subliminal shift of values towards the pathophysiological direction. So far, several MIES symptoms may have been misinterpreted as long COVID symptoms. In any case, the possible MIES triggered by masks contrasts with the WHO definition of health. The exact threshold of harmless and non-pathogenic time wearing a mask should exclusively be determined by further intensive research and studies. Due to the ultimate lack of exclusion of the harmfulness of mask wearing, mask use by the general public should be discouraged.

From the above facts, we conclude that a mask requirement must be reconsidered in a strictly scientific way without any political interference as well as from a humanitarian and ethical point of view. There is an urgent need to balance adverse mask effects with their anticipated efficacy against viral transmission.

**Acknowledgements:** We thank Mrs. Bonita Blankart for proofreading the manuscript.

**Author Contributions:** Conceptualization, K.K., A.S. and O.H. ; Methodology, K.K., A.S. and O.H.; Software, K.K., O.H.; Formal analysis, K.K., O.H. S.W., B.W., S.F., A.P., B.K., S.K.M., P.S. and A.S.; Investigation, K.K., O.H., S.W., B.W., P.S. and A.S.; Physio-metabolic and clinical interpretations, K.K., S.W., S.F., B.K., A.P., P.S. and A.S.; Writing—original draft preparation, K.K., O.H., P.S. and A.S.; Writing—review and editing, K.K., O.H., S.W., B.W., S.F., A.P., B.K., P.S., S.K.M. and A.S.. All authors have read and agreed to the published version of the manuscript  
Correspondence and requests should be addressed to K.K. and P.S..

**Conflict of Interest:** The authors declare no conflict of interest.

**Funding:** This research received no external funding.

#### Literature:

1. Chu DK, Akl EA, Duda S, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet*. 2020;395(10242):1973-1987. doi:10.1016/S0140-6736(20)31142-9
2. Coclite D, Napoletano A, Gianola S, et al. Face Mask Use in the Community for Reducing the Spread of COVID-19: A Systematic Review. *Frontiers in Medicine*. 2021;7. Accessed September 24, 2022. <https://www.frontiersin.org/articles/10.3389/fmed.2020.594269>
3. Howard J, Huang A, Li Z, et al. An evidence review of face masks against COVID-19. *PNAS*. 2021;118(4). doi:10.1073/pnas.2014564118
4. Boretti A. Efficacy of Generalized Face Masking Mandates. *Health Services Research and Managerial Epidemiology*. 2021;8:23333928211058024. doi:10.1177/23333928211058023
5. Fögen Z. The Foegen effect: A mechanism by which facemasks contribute to the COVID-19 case fatality rate. *Medicine (Baltimore)*. 2022;101(7):e28924. doi:10.1097/MD.00000000000028924
6. Janicijevic D, Redondo B, Jiménez R, Lacorzana J, García-Ramos A, Vera J. Intraocular pressure responses to walking with surgical and FFP2/N95 face masks in primary open-angle glaucoma patients. *Graefes Arch Clin Exp Ophthalmol*. 2021;259(8):2373-2378. doi:10.1007/s00417-021-05159-3
7. Keng BMH, Gan WH, Tam YC, Oh CC. Personal protective equipment-related occupational dermatoses during COVID-19 among health care workers: A worldwide systematic review. *JAAD International*. 2021;5:85-95. doi:10.1016/j.jdin.2021.08.004

8. Kisielinski K, Giboni P, Prescher A, et al. Is a Mask That Covers the Mouth and Nose Free from Undesirable Side Effects in Everyday Use and Free of Potential Hazards? *International Journal of Environmental Research and Public Health*. 2021;18(8):4344. doi:10.3390/ijerph18084344
9. Law CSW, Lan PS, Glover GH. Effect of wearing a face mask on fMRI BOLD contrast. *NeuroImage*. 2021;229:117752. doi:10.1016/j.neuroimage.2021.117752
10. Spira B. Correlation Between Mask Compliance and COVID-19 Outcomes in Europe. *Cureus*. 2022;14(4):e24268. doi:10.7759/cureus.24268
11. Vakharia RJ, Jani I, Yadav S, Kurian T. To Study Acute Changes in Brain Oxygenation on MRI in Healthcare Workers Using N95 Mask and PPE Kits for Six Hours a Day. *Indian J Radiol Imaging*. 2021;31(4):893-900. doi:10.1055/s-0041-1741086
12. Sukul P, Schubert JK, Oertel P, et al. FEV manoeuvre induced changes in breath VOC compositions: an unconventional view on lung function tests. *Sci Rep*. 2016;6(1):28029. doi:10.1038/srep28029
13. Sukul P, Schubert JK, Zanaty K, et al. Exhaled breath compositions under varying respiratory rhythms reflects ventilatory variations: translating breathomics towards respiratory medicine. *Scientific Reports*. 2020;10(1):14109. doi:10.1038/s41598-020-70993-0
14. Sukul P, Richter A, Schubert JK, Miekisch W. Deficiency and absence of endogenous isoprene in adults, disqualified its putative origin. *Heliyon*. 2021;7(1). doi:10.1016/j.heliyon.2021.e05922
15. Sukul P, Grzegorzewski S, Broderius C, et al. Physiological and metabolic effects of healthy female aging on exhaled breath biomarkers. *iScience*. 2022;25(2). doi:10.1016/j.isci.2022.103739
16. Remy R, Kemnitz N, Trefz P, et al. Profiling of exhaled volatile organics in the screening scenario of a COVID-19 test center. *iScience*. 2022;25(10). doi:10.1016/j.isci.2022.105195
17. Trefz P, Schmidt SC, Sukul P, Schubert JK, Miekisch W, Fischer DC. Non-Invasive Assessment of Metabolic Adaptation in Paediatric Patients Suffering from Type 1 Diabetes Mellitus. *Journal of Clinical Medicine*. 2019;8(11):1797. doi:10.3390/jcm8111797
18. Sukul P, Schubert JK, Trefz P, Miekisch W. Natural menstrual rhythm and oral contraception diversely affect exhaled breath compositions. *Sci Rep*. 2018;8(1):10838. doi:10.1038/s41598-018-29221-z
19. Löser B, Grabenschröer A, Pugliese G, et al. Changes of Exhaled Volatile Organic Compounds in Postoperative Patients Undergoing Analgesic Treatment: A Prospective Observational Study. *Metabolites*. 2020;10(8):321. doi:10.3390/metabo10080321
20. Sukul P, Bartels J, Fuchs P, et al. Effects of COVID-19 protective face masks and wearing durations on respiratory haemodynamic physiology and exhaled breath constituents. *European Respiratory Journal*. 2022;60(3). doi:10.1183/13993003.00009-2022
21. Engeroff T, Groneberg DA, Niederer D. The Impact of Ubiquitous Face Masks and Filtering Face Piece Application During Rest, Work and Exercise on Gas Exchange, Pulmonary

Function and Physical Performance: A Systematic Review with Meta-analysis. *Sports Medicine - Open*. 2021;7(1):92. doi:10.1186/s40798-021-00388-6

22. Litwinowicz K, Choroszy M, Ornat M, Wróbel A, Waszczuk E. Bayesian network meta-analysis of face masks' impact on human physiology. *Sci Rep*. 2022;12(1):5823. doi:10.1038/s41598-022-09747-z
23. Shaw KA, Zello GA, Butcher SJ, Ko JB, Bertrand L, Chilibeck PD. The impact of face masks on performance and physiological outcomes during exercise: a systematic review and meta-analysis. *Appl Physiol Nutr Metab*. 2021;46(7):693-703. doi:10.1139/apnm-2021-0143
24. Zheng C, Poon ETC, Wan K, Dai Z, Wong SHS. Effects of Wearing a Mask During Exercise on Physiological and Psychological Outcomes in Healthy Individuals: A Systematic Review and Meta-Analysis. *Sports Med*. Published online August 24, 2022. doi:10.1007/s40279-022-01746-4
25. Kunstler B, Newton S, Hill H, et al. P2/N95 respirators & surgical masks to prevent SARS-CoV-2 infection: Effectiveness & adverse effects. *Infect Dis Health*. 2022;27(2):81-95. doi:10.1016/j.idh.2022.01.001
26. Asín-Izquierdo I, Ruiz-Ranz E, Arévalo-Baeza M. The Physiological Effects of Face Masks During Exercise Worn Due to COVID-19: A Systematic Review. *Sports Health*. 2022;14(5):648-655. doi:10.1177/19417381221084661
27. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*. 2015;350:g7647. doi:10.1136/bmj.g7647
28. Huang X, Lin J, Demner-Fushman D. Evaluation of PICO as a knowledge representation for clinical questions. *AMIA Annu Symp Proc*. Published online 2006:359-363.
29. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Systematic Reviews*. 2016;5(1):210. doi:10.1186/s13643-016-0384-4
30. Harrer M, Cuijpers P, Furukawa T, Ebert D. *Doing Meta-Analysis with R: A Hands-On Guide*. Chapman and Hall/CRC; 2021. doi:10.1201/9781003107347
31. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008. doi:10.1136/bmj.j4008
32. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:l4898. doi:10.1136/bmj.l4898
33. casp. Home. CASP - Critical Appraisal Skills Programme. Accessed September 24, 2022. <https://casp-uk.net/>
34. Tufanaru C, Munn Z, Stephenson M, Aromataris E. Fixed or random effects meta-analysis? Common methodological issues in systematic reviews of effectiveness. *Int J Evid Based Healthc*. 2015;13(3):196-207. doi:10.1097/XEB.0000000000000065

35. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials*. 1986;7(3):177-188. doi:10.1016/0197-2456(86)90046-2
36. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*. 2002;21(11):1539-1558. doi:10.1002/sim.1186
37. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. doi:10.1136/bmj.315.7109.629
38. Razi E, Moosavi GA, Omidi K, Khakpour Saebi A, Razi A. Correlation of end-tidal carbon dioxide with arterial carbon dioxide in mechanically ventilated patients. *Arch Trauma Res*. 2012;1(2):58-62. doi:10.5812/atr.6444
39. Contini M, Angelucci A, Aliverti A, et al. Comparison between PtCO<sub>2</sub> and PaCO<sub>2</sub> and Derived Parameters in Heart Failure Patients during Exercise: A Preliminary Study. *Sensors*. 2021;21(19):6666. doi:10.3390/s21196666
40. Górska K, Korczyński P, Maskey-Warzęchowska M, Chazan R, Krenke R. Variability of Transcutaneous Oxygen and Carbon Dioxide Pressure Measurements Associated with Sensor Location. *Adv Exp Med Biol*. 2015;858:39-46. doi:10.1007/5584\_2015\_126
41. Hua W, Zuo Y, Wan R, et al. Short-term skin reactions following use of N95 respirators and medical masks. *Contact Dermatitis*. 2020;83(2):115-121. doi:10.1111/cod.13601
42. Dekkers OM, Vandembroucke JP, Cevallos M, Renehan AG, Altman DG, Egger M. COSMOS-E: Guidance on conducting systematic reviews and meta-analyses of observational studies of etiology. *PLOS Medicine*. 2019;16(2):e1002742. doi:10.1371/journal.pmed.1002742
43. Epstein D, Korytny A, Isenberg Y, et al. Return to training in the COVID-19 era: The physiological effects of face masks during exercise. *Scandinavian Journal of Medicine & Science in Sports*. 2020;n/a(n/a). doi:10.1111/sms.13832
44. Roberge RJ, Kim JH, Powell JB. N95 respirator use during advanced pregnancy. *Am J Infect Control*. 2014;42(10):1097-1100. doi:10.1016/j.ajic.2014.06.025
45. Fikenzer S, Uhe T, Lavall D, et al. Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity. *Clin Res Cardiol*. Published online July 6, 2020:1-9. doi:10.1007/s00392-020-01704-y
46. Lee HP, Wang DY. Objective Assessment of Increase in Breathing Resistance of N95 Respirators on Human Subjects. *Ann Occup Hyg*. 2011;55(8):917-921. doi:10.1093/annhyg/mer065
47. Li Y, Tokura H, Guo YP, et al. Effects of wearing N95 and surgical facemasks on heart rate, thermal stress and subjective sensations. *Int Arch Occup Environ Health*. 2005;78(6):501-509. doi:10.1007/s00420-004-0584-4
48. Mapelli M, Salvioni E, Martino FD, et al. "You can leave your mask on": effects on cardiopulmonary parameters of different airway protection masks at rest and during maximal exercise. *European Respiratory Journal*. Published online January 1, 2021. doi:10.1183/13993003.04473-2020

49. Roberge RJ, Coca A, Williams WJ, Powell JB, Palmiero AJ. Physiological impact of the N95 filtering facepiece respirator on healthcare workers. *Respir Care*. 2010;55(5):569-577.
50. Bharatendu C, Ong JJY, Goh Y, et al. Powered Air Purifying Respirator (PAPR) restores the N95 face mask induced cerebral hemodynamic alterations among Healthcare Workers during COVID-19 Outbreak. *J Neurol Sci*. 2020;417:117078. doi:10.1016/j.jns.2020.117078
51. Butz U. *Rückatmung von Kohlendioxid bei Verwendung von Operationsmasken als hygienischer Mundschutz an medizinischem Fachpersonal*. Universitätsbibliothek der Technischen Universität München; 2005.
52. Dirol H, Alkan E, Sindel M, Ozdemir T, Erbas D. The physiological and disturbing effects of surgical face masks in the COVID-19 era. *BLL*. 2021;122(11):821-825. doi:10.4149/BLL\_2021\_131
53. Georgi C, Haase-Fielitz A, Meretz D, Gäsert L, Butter C. The impact of commonly-worn face masks on physiological parameters and on discomfort during standard work-related physical effort (02.10.2020). *Deutsches Ärzteblatt*. Accessed November 9, 2020. <https://www.aerzteblatt.de/int/archive/article?id=215616>
54. Goh DYT, Mun MW, Lee WLJ, Teoh OH, Rajgor DD. A randomised clinical trial to evaluate the safety, fit, comfort of a novel N95 mask in children. *Scientific Reports*. 2019;9(1):18952. doi:10.1038/s41598-019-55451-w
55. Kim JH, Wu T, Powell JB, Roberge RJ. Physiologic and fit factor profiles of N95 and P100 filtering facepiece respirators for use in hot, humid environments. *Am J Infect Control*. 2016;44(2):194-198. doi:10.1016/j.ajic.2015.08.027
56. Kim JH, Roberge RJ, Powell JB. Effect of external airflow resistive load on postural and exercise-associated cardiovascular and pulmonary responses in pregnancy: a case control study. *BMC Pregnancy and Childbirth*. 2015;15(1):45. doi:10.1186/s12884-015-0474-7
57. Kim JH, Benson SM, Roberge RJ. Pulmonary and heart rate responses to wearing N95 filtering facepiece respirators. *Am J Infect Control*. 2013;41(1):24-27. doi:10.1016/j.ajic.2012.02.037
58. Kyung SY, Kim Y, Hwang H, Park JW, Jeong SH. Risks of N95 Face Mask Use in Subjects With COPD. *Respir Care*. 2020;65(5):658-664. doi:10.4187/respcare.06713
59. Mo Y. Risk and impact of using mask on COPD patients with acute exacerbation during the COVID-19 outbreak: a retrospective study. Published online July 16, 2020. doi:10.21203/rs.3.rs-39747/v1
60. Pifarré F, Zabala DD, Grazioli G, de Yzaguirre i Maura I. COVID 19 and mask in sports. *Apunts Sports Medicine*. Published online June 6, 2020. doi:10.1016/j.apunsm.2020.06.002
61. Rebmann T, Carrico R, Wang J. Physiologic and other effects and compliance with long-term respirator use among medical intensive care unit nurses. *Am J Infect Control*. 2013;41(12):1218-1223. doi:10.1016/j.ajic.2013.02.017

62. Roberge RJ, Kim JH, Benson SM. Absence of consequential changes in physiological, thermal and subjective responses from wearing a surgical mask. *Respiratory Physiology & Neurobiology*. 2012;181(1):29-35. doi:10.1016/j.resp.2012.01.010
63. Tong PSY, Kale AS, Ng K, et al. Respiratory consequences of N95-type Mask usage in pregnant healthcare workers—a controlled clinical study. *Antimicrobial Resistance & Infection Control*. 2015;4(1):48. doi:10.1186/s13756-015-0086-z
64. Zhang G, Li M, Zheng M, et al. Effect of Surgical Masks on Cardiopulmonary Function in Healthy Young Subjects: A Crossover Study. *Front Physiol*. 2021;12:710573. doi:10.3389/fphys.2021.710573
65. Beder A, Büyükköçak U, Sabuncuoğlu H, Keskil ZA, Keskil S. Preliminary report on surgical mask induced deoxygenation during major surgery. *Neurocirugia (Astur)*. 2008;19(2):121-126. doi:10.1016/s1130-1473(08)70235-5
66. Kao TW, Huang KC, Huang YL, Tsai TJ, Hsieh BS, Wu MS. The physiological impact of wearing an N95 mask during hemodialysis as a precaution against SARS in patients with end-stage renal disease. *J Formos Med Assoc*. 2004;103(8):624-628.
67. Toprak E, Bulut AN. The effect of mask use on maternal oxygen saturation in term pregnancies during the COVID-19 process. *Journal of Perinatal Medicine*. 2021;49(2):148-152. doi:10.1515/jpm-2020-0422
68. Tornero-Aguilera JF, Clemente-Suárez VJ. Cognitive and psychophysiological impact of surgical mask use during university lessons. *Physiol Behav*. Published online 2021:113342-113342.
69. Wong AYY, Ling SKK, Louie LHT, et al. Impact of the COVID-19 pandemic on sports and exercise. *Asia Pac J Sports Med Arthrosc Rehabil Technol*. 2020;22:39-44. doi:10.1016/j.asmart.2020.07.006
70. Choudhury A, Singh M, Khurana DK, et al. Physiological Effects of N95 FFP and PPE in Healthcare Workers in COVID Intensive Care Unit: A Prospective Cohort Study. *Indian J Crit Care Med*. 2020;24(12):1169-1173. doi:10.5005/jp-journals-10071-23671
71. Matusiak Ł, Szepietowska M, Krajewski P, Białynicki-Birula R, Szepietowski JC. Inconveniences due to the use of face masks during the COVID-19 pandemic: A survey study of 876 young people. *Dermatologic Therapy*. 2020;33(4):e13567. doi:10.1111/dth.13567
72. Prousa D. Studie zu psychischen und psychovegetativen Beschwerden mit den aktuellen Mund-Nasenschutz-Verordnungen. Published online 2020. doi:10.23668/psycharchives.3135
73. Scarano A, Inchingolo F, Lorusso F. Facial Skin Temperature and Discomfort When Wearing Protective Face Masks: Thermal Infrared Imaging Evaluation and Hands Moving the Mask. *Int J Environ Res Public Health*. 2020;17(13). doi:10.3390/ijerph17134624
74. Jafari E, Togha M, Kazemizadeh H, et al. Evaluation of headache associated with personal protective equipment during COVID-19. *Brain and Behavior*. 2021;11(12):e2435. doi:10.1002/brb3.2435

75. Lan J, Song Z, Miao X, et al. Skin damage among health care workers managing coronavirus disease-2019. *J Am Acad Dermatol*. 2020;82(5):1215-1216. doi:10.1016/j.jaad.2020.03.014
76. Ong JJY, Bharatendu C, Goh Y, et al. Headaches Associated With Personal Protective Equipment - A Cross-Sectional Study Among Frontline Healthcare Workers During COVID-19. *Headache*. 2020;60(5):864-877. doi:10.1111/head.13811
77. Ramirez-Moreno JM. Mask-associated de novo headache in healthcare workers during the Covid-19 pandemic. | medRxiv. Published online 2020. doi:https://doi.org/10.1101/2020.08.07.20167957
78. Rosner E. Adverse Effects of Prolonged Mask Use among Healthcare Professionals during COVID-19. Published online 2020. doi:10.23937/2474-3658/1510130
79. Foo CCI, Goon ATJ, Leow Y, Goh C. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome – a descriptive study in Singapore. *Contact Dermatitis*. 2006;55(5):291-294. doi:10.1111/j.1600-0536.2006.00953.x
80. Klimek L, Huppertz T, Alali A, et al. A new form of irritant rhinitis to filtering facepiece particle (FFP) masks (FFP2/N95/KN95 respirators) during COVID-19 pandemic. *World Allergy Organ J*. 2020;13(10):100474. doi:10.1016/j.waojou.2020.100474
81. Szepietowski JC, Matusiak Ł, Szepietowska M, Krajewski PK, Białynicki-Birula R. Face Mask-induced Itch: A Self-questionnaire Study of 2,315 Responders During the COVID-19 Pandemic. *Acta Derm Venereol*. 2020;100(10):adv00152. doi:10.2340/00015555-3536
82. Techasatian L, Lebsing S, Uppala R, et al. The Effects of the Face Mask on the Skin Underneath: A Prospective Survey During the COVID-19 Pandemic. *J Prim Care Community Health*. 2020;11:2150132720966167. doi:10.1177/2150132720966167
83. Foo CCI, Goon ATJ, Leow Y, Goh C. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome – a descriptive study in Singapore. *Contact Dermatitis*. 2006;55(5):291-294. doi:10.1111/j.1600-0536.2006.00953.x
84. Luckman A, Zeitoun H, Isoni A, et al. *Risk Compensation during COVID-19: The Impact of Face Mask Usage on Social Distancing*. OSF Preprints; 2020. doi:10.31219/osf.io/rb8he
85. Szczesniak D, Ciulkowicz M, Maciaszek J, et al. Psychopathological responses and face mask restrictions during the COVID-19 outbreak: Results from a nationwide survey. *Brain Behav Immun*. 2020;87:161-162. doi:10.1016/j.bbi.2020.05.027
86. Shenal BV, Radonovich LJ, Cheng J, Hodgson M, Bender BS. Discomfort and Exertion Associated with Prolonged Wear of Respiratory Protection in a Health Care Setting. *J Occup Environ Hyg*. 2011;9(1):59-64. doi:10.1080/15459624.2012.635133
87. Park SR, Han J, Yeon YM, Kang NY, Kim E. Effect of face mask on skin characteristics changes during the COVID-19 pandemic. *Skin Res Technol*. 2021;27(4):554-559. doi:10.1111/srt.12983
88. Naylor G, Burke LA, Holman JA. Covid-19 Lockdown Affects Hearing Disability and Handicap in Diverse Ways: A Rapid Online Survey Study. *Ear Hear*. 2020;41(6):1442-1449. doi:10.1097/aud.0000000000000948

89. Thomas F, Allen C, Butts W, Rhoades C, Brandon C, Handrahan DL. Does wearing a surgical facemask or N95-respirator impair radio communication? *Air Med J*. 2011;30(2):97-102. doi:10.1016/j.amj.2010.12.007
90. Heider CA, Álvarez ML, Fuentes-López E, et al. Prevalence of Voice Disorders in Healthcare Workers in the Universal Masking COVID-19 Era. *The Laryngoscope*. 2020;n/a(n/a). doi:10.1002/lary.29172
91. Forgie SE, Reitsma J, Spady D, Wright B, Stobart K. The “fear factor” for surgical masks and face shields, as perceived by children and their parents. *Pediatrics*. 2009;124(4):e777-781. doi:10.1542/peds.2008-3709
92. Islam SR, Prusty D, Dutta R, Maiti S, Chattopadhyay P, Manna SK. Effect of short-term use of FFP2 mask on salivary metabolome of young healthy volunteers: A pilot study. Published online 2022. doi:10.21203/rs.3.rs-1956968/v1
93. Bertoli S, Leone A, De Amicis R, Foppiani A, Osio D, Battezzati A. Effects of wearing a FFP2 mask on indirect calorimetry measurements: A pilot study. *Clin Nutr ESPEN*. 2021;41:443-446. doi:10.1016/j.clnesp.2020.10.015
94. Coniam D. The Impact of Wearing a Face Mask in a High-Stakes Oral Examination: An Exploratory Post-SARS Study in Hong Kong. *Language Assessment Quarterly*. 2005;2(4):235-261. doi:10.1207/s15434311laq0204\_1
95. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Routledge; 1988. doi:10.4324/9780203771587
96. Johnson AT, Scott WH, Lausted CG, Coyne KM, Sahota MS, Johnson MM. Effect of External Dead Volume on Performance While Wearing a Respirator. *AIHAJ - American Industrial Hygiene Association*. 2000;61(5):678-684. doi:10.1080/15298660008984577
97. Xu M, Lei Z, Yang J. Estimating the Dead Space Volume Between a Headform and N95 Filtering Facepiece Respirator Using Microsoft Kinect. *Journal of occupational and environmental hygiene*. 2015;12. doi:10.1080/15459624.2015.1019078
98. Fantin R. The effect of wearing an FFP3 mask (3M TM Aura TM ) with an exhalation valve on gas exchange in medical staff. *Int J Occup Med Environ Health*. Published online April 23, 2021. doi:10.13075/ijomeh.1896.01809
99. Shehade H, Acolty V, Moser M, Oldenhove G. Cutting Edge: Hypoxia-Inducible Factor 1 Negatively Regulates Th1 Function. *The Journal of Immunology*. Published online July 15, 2015. doi:10.4049/jimmunol.1402552
100. Westendorf AM, Skibbe K, Adamczyk A, et al. Hypoxia Enhances Immunosuppression by Inhibiting CD4+ Effector T Cell Function and Promoting Treg Activity. *CPB*. 2017;41(4):1271-1284. doi:10.1159/000464429
101. Sceneay J, Parker BS, Smyth MJ, Möller A. Hypoxia-driven immunosuppression contributes to the pre-metastatic niche. *OncoImmunology*. 2013;2(1):e22355. doi:10.4161/onci.22355

102. Serebrovska ZO, Chong EY, Serebrovska TV, Tumanovska LV, Xi L. Hypoxia, HIF-1 $\alpha$ , and COVID-19: from pathogenic factors to potential therapeutic targets. *Acta Pharmacol Sin.* 2020;41(12):1539-1546. doi:10.1038/s41401-020-00554-8
103. Sturrock A, Zimmerman E, Helms M, Liou TG, Paine III R. Hypoxia induces expression of angiotensin-converting enzyme II in alveolar epithelial cells: Implications for the pathogenesis of acute lung injury in COVID-19. *Physiological Reports.* 2021;9(9):e14854. doi:10.14814/phy2.14854
104. Wang K, Chen W, Zhang Z, et al. CD147-spike protein is a novel route for SARS-CoV-2 infection to host cells. *Signal Transduct Target Ther.* 2020;5(1):283. doi:10.1038/s41392-020-00426-x
105. Chen Z, Mi L, Xu J, et al. Function of HAb18G/CD147 in invasion of host cells by severe acute respiratory syndrome coronavirus. *J Infect Dis.* 2005;191(5):755-760. doi:10.1086/427811
106. Ke X, Fei F, Chen Y, et al. Hypoxia upregulates CD147 through a combined effect of HIF-1 $\alpha$  and Sp1 to promote glycolysis and tumor progression in epithelial solid tumors. *Carcinogenesis.* 2012;33(8):1598-1607. doi:10.1093/carcin/bgs196
107. Aggarwal BB. Nuclear factor-kappaB: the enemy within. *Cancer Cell.* 2004;6(3):203-208. doi:10.1016/j.ccr.2004.09.003
108. Blylock RL. Immunoexcitatory mechanisms in glioma proliferation, invasion and occasional metastasis. *Surg Neurol Int.* 2013;4:15. doi:10.4103/2152-7806.106577
109. Li Y, Xu J, Chen L, et al. HAb18G (CD147), a cancer-associated biomarker and its role in cancer detection. *Histopathology.* 2009;54(6):677-687. doi:10.1111/j.1365-2559.2009.03280.x
110. Walach H, Traindl H, Prentice J, et al. Carbon dioxide rises beyond acceptable safety levels in children under nose and mouth covering: Results of an experimental measurement study in healthy children. *Environmental Research.* 2022;212:113564. doi:10.1016/j.envres.2022.113564
111. Acuti Martellucci C, Flacco ME, Martellucci M, Violante FS, Manzoli L. Inhaled CO<sub>2</sub> Concentration While Wearing Face Masks: A Pilot Study Using Capnography. *Environ Health Insights.* 2022;16:11786302221123572. doi:10.1177/11786302221123573
112. Hoffman WE, Charbel FT, Edelman G, Ausman JI. Brain tissue acid-base response to hypercapnia in neurosurgical patients. *Neurol Res.* 1995;17(6):417-420.
113. Huo X long, Min J jing, Pan C yu, et al. Efficacy of Lovastatin on Learning and Memory Deficits Caused by Chronic Intermittent Hypoxia-Hypercapnia: Through Regulation of NR2B-Containing NMDA Receptor-ERK Pathway. *PLOS ONE.* 2014;9(4):e94278. doi:10.1371/journal.pone.0094278
114. Sikter A, Faludi G, Rihmer Z. The role of carbon dioxide (and intracellular pH) in the pathomechanism of several mental disorders. Are the diseases of civilization caused by learnt behaviour, not the stress itself? *Neuropsychopharmacol Hung.* 2009;11(3):161-173.

115. Schaefer KE. Respiratory adaptation to chronic hypercapnia. *Ann N Y Acad Sci.* 1963;109:772-782. doi:10.1111/j.1749-6632.1963.tb13505.x
116. Schaefer KE, Douglas WH, Messier AA, Shea ML, Gohman PA. Effect of prolonged exposure to 0.5% CO<sub>2</sub> on kidney calcification and ultrastructure of lungs. *Undersea Biomed Res.* 1979;6 Suppl:S155-161.
117. Benjamin Ezraty, Maïalène Chabalier, Adrien Ducret, Etienne Maisonneuve, Sam Dukan. CO<sub>2</sub> exacerbates oxygen toxicity. *EMBO reports.* 2011;12(4):321-326. doi:10.1038/embor.2011.7
118. Guais A, Brand G, Jacquot L, et al. Toxicity of Carbon Dioxide: A Review. *Chem Res Toxicol.* 2011;24(12):2061-2070. doi:10.1021/tx200220r
119. Beheshti A, Cekanaviciute E, Smith DJ, Costes SV. Global transcriptomic analysis suggests carbon dioxide as an environmental stressor in spaceflight: A systems biology GeneLab case study. *Sci Rep.* 2018;8(1):4191. doi:10.1038/s41598-018-22613-1
120. Jacobson TA, Kler JS, Hernke MT, Braun RK, Meyer KC, Funk WE. Direct human health risks of increased atmospheric carbon dioxide. *Nat Sustain.* 2019;2(8):691-701. doi:10.1038/s41893-019-0323-1
121. Thom SR, Bhopale VM, Hu J, Yang M. Inflammatory responses to acute elevations of carbon dioxide in mice. *Journal of Applied Physiology.* 2017;123(2):297-302. doi:10.1152/jappphysiol.00343.2017
122. Zappulla D. Environmental Stress, Erythrocyte Dysfunctions, Inflammation, and the Metabolic Syndrome: Adaptations to CO<sub>2</sub> Increases? *Journal of the CardioMetabolic Syndrome.* 2008;3(1):30-34. doi:10.1111/j.1559-4572.2008.07263.x
123. Forrest VJ, Kang YH, McClain DE, Robinson DH, Ramakrishnan N. Oxidative stress-induced apoptosis prevented by Trolox. *Free Radic Biol Med.* 1994;16(6):675-684. doi:10.1016/0891-5849(94)90182-1
124. Veselá A, Wilhelm J. The role of carbon dioxide in free radical reactions of the organism. *Physiol Res.* 2002;51(4):335-339.
125. Sukul P, Trefz P, Kamysek S, Schubert JK, Miekisch W. Instant effects of changing body positions on compositions of exhaled breath. *J Breath Res.* 2015;9(4):047105. doi:10.1088/1752-7155/9/4/047105
126. Xie A, Skatrud JB, Puleo DS, Morgan BJ. Exposure to hypoxia produces long-lasting sympathetic activation in humans. *J Appl Physiol (1985).* 2001;91(4):1555-1562. doi:10.1152/jappl.2001.91.4.1555
127. Delanghe L, Cauwenberghs E, Spacova I, et al. Cotton and Surgical Face Masks in Community Settings: Bacterial Contamination and Face Mask Hygiene. *Frontiers in Medicine.* 2021;8. Accessed September 24, 2022. <https://www.frontiersin.org/articles/10.3389/fmed.2021.732047>
128. Kisielinski K, Wojtasik B, Suitability of Rose Bengal sodium salt staining for visualisation of face mask contamination by living organisms. *AIMSES.* 2022;9(2):218-231. doi:10.3934/environsci.2022015


129. Park AM, Khadka S, Sato F, et al. Bacterial and fungal isolation from face masks under the COVID-19 pandemic. *Sci Rep*. 2022;12(1):11361. doi:10.1038/s41598-022-15409-x
130. Arora U, Priyadarshi M, Katiyar V, et al. Risk factors for Coronavirus disease-associated mucormycosis. *J Infect*. 2022;84(3):383-390. doi:10.1016/j.jinf.2021.12.039
131. Ellingsen I, Sydnnes G, Hauge A, Zwart JA, Liestøl K, Nicolaysen G. CO<sub>2</sub> sensitivity in humans breathing 1 or 2% CO<sub>2</sub> in air. *Acta Physiologica Scandinavica*. 1987;129(2):195-202. doi:10.1111/j.1748-1716.1987.tb08059.x
132. Langford NJ. Carbon Dioxide Poisoning. *Toxicol Rev*. 2005;24(4):229-235. doi:10.2165/00139709-200524040-00003
133. Kisielinski K, Wagner S, Hirsch O, Klosterhalfen B, Prescher A. *Possible Toxicity of Chronic Carbon Dioxide Exposure Associated with Mask Use, Particularly in Pregnant Women, Children and Adolescents -a Scoping Review*. Preprints; 2022. doi:10.22541/au.164269384.47200789/v1
134. Grollman A. Physiological variations in the cardiac output of man. *American Journal of Physiology-Legacy Content*. 1929;89(3):584-588. doi:10.1152/ajplegacy.1929.89.3.584
135. Kiely DG, Cargill RI, Lipworth BJ. Effects of hypercapnia on hemodynamic, inotropic, lusitropic, and electrophysiologic indices in humans. *Chest*. 1996;109(5):1215-1221. doi:10.1378/chest.109.5.1215
136. Zheng G qing, Wang Y, Wang X tong. Chronic hypoxia-hypercapnia influences cognitive function: a possible new model of cognitive dysfunction in chronic obstructive pulmonary disease. *Med Hypotheses*. 2008;71(1):111-113. doi:10.1016/j.mehy.2008.01.025
137. Loeb M, Bartholomew A, Hashmi M, et al. Medical Masks Versus N95 Respirators for Preventing COVID-19 Among Health Care Workers. *Ann Intern Med*. Published online November 29, 2022. doi:10.7326/M22-1966
138. Hua D, Xu Y, Heiduschka P, et al. Retina Vascular Perfusion Dynamics During Exercise With and Without Face Masks in Healthy Young Adults: An OCT Angiography Study. *Translational Vision Science & Technology*. 2021;10(3):23. doi:10.1167/tvst.10.3.23
139. D'Souza S, Vaidya T, Nair AP, et al. Altered Ocular Surface Health Status and Tear Film Immune Profile Due to Prolonged Daily Mask Wear in Health Care Workers. *Biomedicines*. 2022;10(5):1160. doi:10.3390/biomedicines10051160
140. Redlich CA, Sparer J, Cullen MR. Sick-building syndrome. *Lancet*. 1997;349(9057):1013-1016. doi:10.1016/S0140-6736(96)07220-0
141. Kim EA, Kang SK. Historical review of the List of Occupational Diseases recommended by the International Labour organization (ILO). *Annals of Occupational and Environmental Medicine*. 2013;25(1):14. doi:10.1186/2052-4374-25-14
142. Micheli LJ, Allison G. Lumbar spine injury in the young athlete. *Rev Bras Med Esporte*. 1999;5:59-65. doi:10.1590/S1517-86921999000200005

143. Van Mechelen M, Lories RJ. Microtrauma: no longer to be ignored in spondyloarthritis? *Curr Opin Rheumatol*. 2016;28(2):176-180. doi:10.1097/BOR.0000000000000254
144. Wilder RP, Sethi S. Overuse injuries: tendinopathies, stress fractures, compartment syndrome, and shin splints. *Clin Sports Med*. 2004;23(1):55-81, vi. doi:10.1016/S0278-5919(03)00085-1
145. Stahl S, Stahl AS, Meisner C, Rahmanian-Schwarz A, Schaller HE, Lotter O. A systematic review of the etiopathogenesis of Kienböck's disease and a critical appraisal of its recognition as an occupational disease related to hand-arm vibration. *BMC Musculoskeletal Disorders*. 2012;13(1):225. doi:10.1186/1471-2474-13-225
146. Cher I. Blink-related microtrauma: when the ocular surface harms itself. *Clinical & Experimental Ophthalmology*. 2003;31(3):183-190. doi:10.1046/j.1442-9071.2003.00633.x
147. Boulanger G. (t)rauma With a Small t? *Contemporary Psychoanalysis*. 2016;52(1):143-149. doi:10.1080/00107530.2016.1149398
148. Crastnopol M. *Micro-Trauma: A Psychoanalytic Understanding of Cumulative Psychic Injury*. Routledge; 2015. doi:10.4324/9780203878897
149. Seides R. Should the current DSM-IV-TR definition for PTSD be expanded to include serial and multiple microtraumas as aetiologies? *J Psychiatr Ment Health Nurs*. 2010;17(8):725-731. doi:10.1111/j.1365-2850.2010.01591.x
150. Kisielinski K, Niedhart C, Schneider U, Niethard FU. Osteonecrosis 15 years after femoral neck fracture and long-term low-dose inhaled corticosteroid therapy. *Joint Bone Spine*. 2004;71(3):237-239. doi:10.1016/S1297-319X(03)00115-5
151. Miravittles M, Auladell-Rispau A, Monteagudo M, et al. Systematic review on long-term adverse effects of inhaled corticosteroids in the treatment of COPD. *European Respiratory Review*. 2021;30(160). doi:10.1183/16000617.0075-2021
152. Houston TP. The silent killer: environmental tobacco smoke. *J Fam Pract*. 1991;32(5):457-458.
153. Shaldon S, Vienken J. Beyond The Current Paradigm: Recent Advances in The Understanding of Sodium Handling – Guest Editors: Stanley Shaldon and Joerg Vienken: Salt, the Neglected Silent Killer. *Seminars in Dialysis*. 2009;22(3):264-266. doi:10.1111/j.1525-139X.2009.00606.x
154. Alasfar RH, Isaifan RJ. Aluminum environmental pollution: the silent killer. *Environ Sci Pollut Res*. 2021;28(33):44587-44597. doi:10.1007/s11356-021-14700-0
155. Nawrot TS, Staessen JA. Low-Level Environmental Exposure to Lead Unmasked as Silent Killer. *Circulation*. 2006;114(13):1347-1349. doi:10.1161/CIRCULATIONAHA.106.650440
156. Zaynab M, Fatima M, Sharif Y, et al. Health and environmental effects of silent killers Organochlorine pesticides and polychlorinated biphenyl. *Journal of King Saud University - Science*. 2021;33(6):101511. doi:10.1016/j.jksus.2021.101511
157. Huckelba AL, Van Lange PAM. The Silent Killer: Consequences of Climate Change and How to Survive Past the Year 2050. *Sustainability*. 2020;12(9):3757. doi:10.3390/su12093757

158. Chen C, Hauptert SR, Zimmermann L, Shi X, Fritsche LG, Mukherjee B. Global Prevalence of Post COVID-19 Condition or Long COVID: A Meta-Analysis and Systematic Review. *J Infect Dis*. Published online April 16, 2022;jiac136. doi:10.1093/infdis/jiac136
159. Conference IH. WHO - Constitution of the World Health Organization. 1946. *Bulletin of the World Health Organization*. 2002;80(12):983-984.
160. Coma E, Català M, Méndez-Boo L, et al. Unravelling the role of the mandatory use of face covering masks for the control of SARS-CoV-2 in schools: a quasi-experimental study nested in a population-based cohort in Catalonia (Spain). *Archives of Disease in Childhood*. Published online August 23, 2022. doi:10.1136/archdischild-2022-324172
161. Fønhus MS, Dalsbø TK, Brurberg KG. *Facemasks to Prevent Transmission of Respiratory Illness, Such as COVID-19*. Norwegian Institute of Public Health; 2021. Accessed November 7, 2021. <https://fhi.brage.unit.no/fhi-xmlui/handle/11250/2756758>
162. Jefferson T, Mar CBD, Dooley L, et al. Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database of Systematic Reviews*. 2020;(11). doi:10.1002/14651858.CD006207.pub5
163. Bundgaard H, Bundgaard JS, Raaschou-Pedersen DET, et al. Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers. *Ann Intern Med*. Published online November 18, 2020. doi:10.7326/M20-6817
164. Abaluck J, Kwong LH, Styczynski A, et al. Impact of community masking on COVID-19: A cluster-randomized trial in Bangladesh. *Science*. 2021;375(6577):eabi9069. doi:10.1126/science.abi9069
165. Glasziou PP, Michie S, Fretheim A. Public health measures for covid-19. *BMJ*. 2021;375:n2729. doi:10.1136/bmj.n2729
166. Wang MX, Gwee SXW, Chua PEY, Pang J. Effectiveness of Surgical Face Masks in Reducing Acute Respiratory Infections in Non-Healthcare Settings: A Systematic Review and Meta-Analysis. *Frontiers in Medicine*. 2020;7. Accessed November 26, 2022. <https://www.frontiersin.org/articles/10.3389/fmed.2020.564280>
167. Xiao J, Shiu EYC, Gao H, et al. Nonpharmaceutical Measures for Pandemic Influenza in Nonhealthcare Settings—Personal Protective and Environmental Measures - Volume 26, Number 5—May 2020 - Emerging Infectious Diseases journal - CDC. doi:10.3201/eid2605.190994
168. Mader S, Rüttenauer T. The Effects of Non-pharmaceutical Interventions on COVID-19 Mortality: A Generalized Synthetic Control Approach Across 169 Countries. *Frontiers in Public Health*. 2022;10. Accessed November 28, 2022. <https://www.frontiersin.org/articles/10.3389/fpubh.2022.820642>
169. Asadi S, Cappa CD, Barreda S, Wexler AS, Bouvier NM, Ristenpart WD. Efficacy of masks and face coverings in controlling outward aerosol particle emission from expiratory activities. *Scientific Reports*. 2020;10(1):15665. doi:10.1038/s41598-020-72798-7

170. Schlegtendal A, Eitner L, Falkenstein M, et al. To Mask or Not to Mask—Evaluation of Cognitive Performance in Children Wearing Face Masks during School Lessons (MasKids). *Children*. 2022;9(1):95. doi:10.3390/children9010095

Figures 1-13:




**Figure 1.** PRISMA flow chart of the systematic review. From initial 2168, fifty-four studies were later included in the qualitative synthesis. Finally 37 studies were evaluated statistically in the meta-analysis (quantitative analysis).


Meta-analysis of biochemical outcomes

A) SpO<sub>2</sub>


Blood oxygen saturation when using a mask (general)



Blood oxygen saturation when using a surgical mask




Blood oxygen saturation when using a N95 mask




B) CO<sub>2</sub> (evaluation of PtCO<sub>2</sub>, ETCO<sub>2</sub> and PaCO<sub>2</sub>)

Carbon dioxide blood content (mmHg) in mask use (general)



Carbon dioxide blood content (mmHg) in surgical mask use



Carbon dioxide blood content (mmHg) in N95 mask use




Figure 2.

Forest (left) and Funnel plots (right) of meta-analysis of blood oxygenation and blood carbon dioxide outcomes while wearing a face mask. All face mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: □=surgical mask ■=N95 mask.



A: Blood oxygen is significantly lowered in mask use. In the subgroup analysis this could also be found for N95 mask use. From the pooled analysis, it seems, that N95 mask may be responsible for a larger SpO<sub>2</sub> drop than surgical masks. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always lower O<sub>2</sub>-values than the surgical masks.

B: In the pooled analysis, blood carbon dioxide (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>) is significantly elevated in mask use. This could be found for general mask use and in the subgroup analysis for surgical mask, and also for N95 mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher CO<sub>2</sub>-values than the surgical masks.



### Meta-analysis of respiratory outcomes

#### A) Ventilation



##### Ventilation (l/min) when using a mask (general)



##### Ventilation (l/min) when using a surgical mask





##### Ventilation (l/min) when using a N95 mask



#### B) Respiratory rate


##### Breathing frequency (breaths/min) when using a mask (general)



**Figure 3.** Forest (left) and Funnel plots (right) of meta-analysis of physiological respiratory outcomes while wearing a face mask. **A** shows results for ventilation ( $V_E$ ), **B** for respiratory rate (RR). All face mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: □=surgical mask ■=N95 mask.

**A:** Breathing volume is significantly lowered in mask use in the pooled analysis. This could be found for general, for surgical, and N95 mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always lower ventilation ( $V_E$ ) than the surgical masks.


**B:** No statistical difference could be found regarding respiratory rate in mask use in the pooled analysis, even in the subgroup analysis (not shown).



**Figure 4.** Forest (left) and Funnel plots (right) of metaanalysis of the physiological cardiovascular outcomes systolic blood pressure (SBP) and heart rate (HR). All controlled intervention studies in which measurements were taken during physical activity with face masks were included (exclusion of rest situation and pre-post studies). All face masks types are initially considered together, later if possible subgroups (surgical and N95) are evaluated. If studies evaluate two different mask types in parallel, the corresponding studies are marked: □=surgical mask ■=N95 mask.

**A:** Systolic blood pressure is elevated in the mask condition and also for the subgroup of surgical mask. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher SBP than the surgical mask, however this effect was not statistically significant.

**B:** For the N95 mask condition a low significance for a slight increase in heart rate could be found. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher HR than the surgical mask, and this effect was statistically significant.



**Figure 5.** Forest (left) and Funnel plots (right) of meta-analysis of physical outcomes while wearing a face mask. **A** shows results for temperature of skin, **B** for air humidity underneath the face mask. All mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: □=surgical mask ■=N95 mask.

**A:** Skin covered by mask has a significantly higher temperature during rest and activity. This could be found for general mask use and for N95 mask use but not for surgical mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded higher temperatures than the surgical mask, but this could not be analysed further due to lack of further studies comparing both conditions.



**B:** The dead space covered by mask has a significantly higher air humidity in the pooled analysis.





Meta-analysis of N95 mask vs surgical mask

A) Biochemical comparison

Blood oxygen saturation (SpO<sub>2</sub>) when using a N95 mask vs surgical mask





Carbon dioxide blood content (mmHg) in N95 mask use vs surgical mask





B) Cardiorespiratory comparison



Ventilation (l/min) when using a N95 mask vs surgical mask



Heart rate (beats/min) when using a N95 mask vs surgical mask





Systolic blood pressure (mmHg) when using a N95 vs surgical mask





C) Measured subjective sensations comparison

Perceived discomfort when using a N95 mask vs surgical mask



Perceived exertion when using a N95 mask vs surgical mask



Perceived humidity when using a N95 mask vs surgical mask






Figure 7.


Results comparing the N95 to the surgical mask in the meta-analysis. Forest (left) and Funnel plots (right) of meta-analysis of diverse outcomes while wearing a N95 mask vs surgical mask are shown. **A** depicts the biochemical, **B** the cardiorespiratory outcomes and **C** the subjective sensations outcomes. N95 mask leads to measurably less favourable results compared to the surgical mask, significantly for oxygenation (decrease), heart rate (increase), discomfort and humidity (both increases). This trend was also evident for minute volume (decrease), CO<sub>2</sub> and systolic blood pressure (both increases), but in those comparisons not statistically significant due to too few includable studies.

### Meta-analysis of pooled symptom prevalence while wearing a face mask


#### Headache prevalence when using a face mask




#### Acne prevalence when using a face mask




#### Skin irritation prevalence when using a face mask




#### Shortness of breath prevalence when using a face mask




#### Heat prevalence when using a face mask




#### Itch prevalence when using a face mask




#### Voice disorder prevalence when using a face mask



#### Dizziness prevalence when using a face mask



**Figure 8.** Forest (left) and Funnel plots (right) of meta-analysis of pooled symptom prevalence while wearing a face mask. Headache (62%), acne (38%), skin irritation (36%), shortness of breath (33%), heat (26%), itch (26%), voice disorder (23%) and dizziness (5%) while wearing a mask are significant in the evaluated population (n=8128).



**Figure 9.** Summary of pooled metaanalytic evaluation of biochemical (A) and physical effects (B) during face mask use. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the confidence intervals.

**A:** For carbon dioxide rise in the blood there is a medium effect size of >0.5 and for oxygen drop a small effect size of >0.2 regarding the standard mean difference values thresholds according to Cohen 1988.

**B:** For elevated Humidity and Temperature rise under the face mask there is a strong effect size of  $\geq 0.8$ .


The metaanalytical statistical data were as follows:

Oxygen (SpO<sub>2</sub>): SMD -0.24, 95% CI -0.38 to -0.11, Z=3.53, p=0.0004;

Carbon dioxide (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>): SMD +0.64, 95% CI 0.31 to 0.96, Z=3.86, p=0.0001;

Humidity: SMD +2.24, 95% CI 1.32 to 3.17, Z=4.75, p<0.00001;

Temperature: SMD +0.8, 95% CI 0.23 to 1.38, Z=2.72, p=0.008.



**Figure 10.** Summary of pooled metaanalytic evaluation of cardiorespiratory effects during face mask use. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the confidence intervals.

Clear effects for a decrease in ventilation and tidal volume are illustrated, no effect for respiratory rate and weak to low effect for increase in heart rate and systolic blood pressure. For ventilation there is a medium effect size of  $>0.5$  with a small effect size of  $>0.2$  for tidal volume of the standard mean difference values according to Cohen 1988.

The metaanalytical statistical data were as follows:

Ventilation: SMD -0.72, 95% CI -0.99 to -0.46,  $Z=5.36$ ,  $p<0.00001$ ;


Tidal volume: SMD -0.37, 95% CI -0.63 to -0.11,  $Z=2.82$ ,  $p=0.005$ ;

Respiratory rate: SMD +0.01, 95% CI -0.29 to 0.30,  $Z=0.08$ ,  $p=0.94$ ;

Heart rate: SMD +0.11, 95% CI -0.05 to 0.28,  $Z=1.34$ ,  $p=0.18$ ;

Systolic blood pressure: SMD +0.17, 95% CI 0.03 to 0.32,  $Z=2.39$ ,  $p=0.02$ .





**Figure 12.** Summary of pooled metaanalytic evaluation of face mask-wearing sensations measured with standardised Borg-, Likert-, VAS-scales or similar. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the confidence intervals.

Five out of 6 complaint categories (83%) are above the strong effect size threshold of >0.8 of the standard mean difference values according to Cohen 1988.

The metaanalytical statistical data were as follows (SMD=standard mean difference):

Itch: SMD +1.57, 95 %CI -0.08 to 3.23, Z=1.86, p=0.06;


Shortness of breath: SMD +1.46, 95% CI 0.42 to 2.50, Z=2.75, p=0.006;

Discomfort: SMD +1.16, 95% CI 0.58 to 1.73, Z=3.94, p<0.0001;

Exertion: SMD +0.9, 95 % CI 0.57 to 1.23, Z=5.31, p<0.00001;

Humidity: SMD +0.9, 95% CI 0.34 to 1.46, Z=3.17, p=0.002;

Heat: SMD +0.77, 95% CI 0.29 to 1.26, Z=3.11, p=0.002.




**Figure 13.** Representation of symptom prevalence in % during face mask use as the area of the circles. Along the X-axis, the main recorded symptoms are listed. The higher the prevalence, the bigger the circles and the more often the symptoms.

The Y-axis gives the probability of non-random occurrence of the symptoms and includes the statistical Z-value. Thus, the higher the circles are arranged, the more robust is the relationship to face mask wearing.

The metaanalytical statistical data were as follows:

- Headache: 62% (95% CI 48-75%), Z=8.77, p<0.00001;
- Acne: 38% (95% CI 22-54%), Z=4.58, p<0.00001;
- Skin irritation: 36% (95% CI 24-49%), Z=5.61, p<0.00001;
- Shortness of breath: 33% (95% CI 23-44%), Z=6.28, p<0.00001;
- Heat: 28% (95% CI 15-43%), Z=4.72, p<0.00001;
- Itch: 26% (95% CI 15-36%), Z=4.77, p<0.00001;
- Voice disorder 23% (95% CI 2-43%), Z=2.15, p<0.03;
- Dizziness 5% (95% CI 1-9%), Z=2.5, p=0.01.

# Figures




**Figure 1**

PRISMA flow chart of the systematic review. From initial 2168, fifty-four studies were later included in the qualitative synthesis. Finally 37 studies were evaluated statistically in the meta-analysis (quantitative analysis).


## Meta-analysis of biochemical outcomes

### A) SpO<sub>2</sub>


#### Blood oxygen saturation when using a mask (general)



#### Blood oxygen saturation when using a surgical mask




#### Blood oxygen saturation when using a N95 mask




### B) CO<sub>2</sub> (evaluation of PTCO<sub>2</sub>, ETCO<sub>2</sub> und PaCO<sub>2</sub>)


#### Carbon dioxide blood content (mmHg) in mask use (general)



#### Carbon dioxide blood content (mmHg) in surgical mask use



#### Carbon dioxide blood content (mmHg) in N95 mask use



## Figure 2

Forest (left) and Funnel plots (right) of meta-analysis of blood oxygenation and blood carbon dioxide outcomes while wearing a face mask. All face mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: =surgical mask =N95 mask.

**A:** Blood oxygen is significantly lowered in mask use. In the subgroup analysis this could also be found for N95 mask use. From the pooled analysis, it seems, that N95 mask may be responsible for a larger SpO<sub>2</sub> drop than surgical masks. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always lower O<sub>2</sub>-values than the surgical masks.

**B:** In the pooled analysis, blood carbon dioxide (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>) is significantly elevated in mask use. This could be found for general mask use and in the subgroup analysis for surgical mask, and also for N95 mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher CO<sub>2</sub>-values than the surgical masks.



# Meta-analysis of respiratory outcomes

## A) Ventilation

### Ventilation (l/min) when using a mask (general)

Study or Subgroup	mask			no mask			Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Fikenzler □ 2020	114	23.3	12	131	27.8	12	10.2%	-0.64 [-1.46, 0.18]
Fikenzler ■ 2020	98.8	18.6	12	131	27.8	12	8.6%	-1.31 [-2.21, -0.42]
Mapelli □ 2021	76.2	21.6	12	92.3	26	12	10.2%	-0.65 [-1.48, 0.17]
Mapelli ■ 2021	71.6	21.2	12	92.3	26	12	9.8%	-0.84 [-1.68, -0.00]
Zhang 2021	55.1	17.3	71	66.5	17.9	71	61.1%	-0.64 [-0.98, -0.31]
<b>Total (95% CI)</b>	<b>119</b>			<b>119</b>			<b>100.0%</b>	<b>-0.72 [-0.99, -0.46]</b>



Heterogeneity: Tau<sup>2</sup> = 0.00; Chi<sup>2</sup> = 2.02, df = 4 (P = 0.73); I<sup>2</sup> = 0%  
Test for overall effect: Z = 5.36 (P < 0.00001)



### Ventilation (l/min) when using a surgical mask

Study or Subgroup	surgical mask			no mask			Weight	Std. Mean Difference IV, Random, 95% CI	Year
	Mean	SD	Total	Mean	SD	Total			
Fikenzler 2020	114	23.3	12	131	27.8	12	12.6%	-0.64 [-1.46, 0.18]	2020
Mapelli 2021	76.2	21.6	12	92.3	26	12	12.5%	-0.65 [-1.48, 0.17]	2021
Zhang 2021	55.1	17.3	71	66.5	17.9	71	74.9%	-0.64 [-0.98, -0.31]	2021
<b>Total (95% CI)</b>	<b>95</b>			<b>95</b>			<b>100.0%</b>	<b>-0.64 [-0.94, -0.35]</b>	



Heterogeneity: Tau<sup>2</sup> = 0.00; Chi<sup>2</sup> = 0.00, df = 2 (P = 1.00); I<sup>2</sup> = 0%  
Test for overall effect: Z = 4.32 (P < 0.0001)



### Ventilation (l/min) when using a N95 mask

Study or Subgroup	N95 mask			no mask			Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Fikenzler 2020	98.8	18.6	12	131	27.8	12	46.8%	-1.31 [-2.21, -0.42]
Mapelli 2021	71.6	21.2	12	92.3	26	12	53.2%	-0.84 [-1.68, -0.00]
<b>Total (95% CI)</b>	<b>24</b>			<b>24</b>			<b>100.0%</b>	<b>-1.06 [-1.68, -0.45]</b>

Heterogeneity: Tau<sup>2</sup> = 0.00; Chi<sup>2</sup> = 0.57, df = 1 (P = 0.45); I<sup>2</sup> = 0%  
Test for overall effect: Z = 3.39 (P = 0.0007)



## B) Respiratory rate

### Breathing frequency (breaths/min) when using a mask (general)

Study or Subgroup	mask			no mask			Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Dirol 2021	19.7	3.21	100	17.46	3.31	100	9.8%	0.68 [0.40, 0.97]
Fikenzler □ 2020	39.3	6.2	12	40.9	5.1	12	5.9%	-0.27 [-1.08, 0.53]
Fikenzler ■ 2020	36.8	5.9	12	40.9	5.1	12	5.7%	-0.72 [-1.55, 0.11]
Georgi □ 2020	28.2	8.5	24	26.4	6.1	24	7.6%	0.24 [-0.33, 0.81]
Georgi ■ 2020	29	9.8	24	26.4	6.1	24	7.6%	0.31 [-0.26, 0.88]
Kim 2013	24.1	3.7	16	21.7	3.4	16	6.5%	0.66 [-0.06, 1.37]
Kim 2016	28.4	3.2	12	28.1	7.1	12	5.9%	0.05 [-0.75, 0.85]
Kyung 2020	25.7	7.5	90	23.3	2.6	90	9.7%	0.43 [0.13, 0.72]
Mapelli □ 2021	37.7	5.5	12	41.5	8	12	5.8%	-0.53 [-1.35, 0.28]
Mapelli ■ 2021	37.1	4.5	12	41.5	8	12	5.8%	-0.65 [-1.48, 0.17]
Roberge 2010	26.6	6.8	10	27.7	8.6	10	5.4%	-0.14 [-1.01, 0.74]
Roberge 2012	24.7	3.7	20	23.7	2.7	20	7.2%	0.30 [-0.32, 0.93]
Roberge 2014	24.9	6.1	22	26.4	4.2	22	7.4%	-0.28 [-0.88, 0.31]
Zhang 2021	33.8	7.98	71	37.9	6.72	71	9.4%	-0.55 [-0.89, -0.22]
<b>Total (95% CI)</b>	<b>437</b>			<b>437</b>			<b>100.0%</b>	<b>0.01 [-0.28, 0.30]</b>

Heterogeneity: Tau<sup>2</sup> = 0.20; Chi<sup>2</sup> = 50.36, df = 13 (P < 0.00001); I<sup>2</sup> = 74%  
Test for overall effect: Z = 0.08 (P = 0.94)





Figure 3

Forest (left) and Funnel plots (right) of meta-analysis of physiological respiratory outcomes while wearing a face mask. **A** shows results for ventilation ( $V_E$ ), **B** for respiratory rate (RR). All face mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: □ =surgical mask ■ =N95 mask.


**A:** Breathing volume is significantly lowered in mask use in the pooled analysis. This could be found for general, for surgical, and N95 mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always lower ventilation ( $V_E$ ) than the surgical masks.

**B:** No statistical difference could be found regarding respiratory rate in mask use in the pooled analysis, even in the subgroup analysis (not shown).


## Meta-analysis of cardiovascular outcomes

### A) Systolic blood pressure (SBP)


#### Systolic blood pressure (mmHg) when using a mask (general)



#### Systolic blood pressure (mmHg) when using a surgical mask




#### Systolic blood pressure (mmHg) when using a N95 mask




### B) Heart rate

#### Heart rate (beats/min) when using a mask (general)



#### Heart rate (beats/min) when using a surgical mask



#### Heart rate (beats/min) when using a N95 mask




Figure 4

Forest (left) and Funnel plots (right) of metaanalysis of the physiological cardiovascular outcomes systolic blood pressure (SBP) and heart rate (HR). All controlled intervention studies in which measurements were taken during physical activity with face masks were included (exclusion of rest situation and pre-post studies). All face masks types are initially considered together, later if possible

subgroups (surgical and N95) are evaluated. If studies evaluate two different mask types in parallel, the corresponding studies are marked: =surgical mask =N95 mask.

**A:** Systolic blood pressure is elevated in the mask condition and also for the subgroup of surgical mask. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher SBP than the surgical mask, however this effect was not statistically significant.

**B:** For the N95 mask condition a low significance for a slight increase in heart rate could be found. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher HR than the surgical mask, and this effect was statistically significant.


# Meta-analysis of physical outcomes

## A) Temperature (skin in °C)

### Skin temperature below a mask during use

Study or Subgroup	mask			no mask			Weight	Std. Mean Difference	
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	95% CI
Kim 2016	35	0.7	12	34.8	0.7	12	18.2%	0.28	[-0.53, 1.08]
Park 2020	35.133	1.229	21	33.5	1.235	21	20.4%	1.30	[0.63, 1.97]
Roberge 2012	33.7	0.88	20	31.94	1.1	20	19.3%	1.73	[0.99, 2.47]
Scarano □ 2020	35.9	3.4	20	35.2	3.1	20	21.2%	0.21	[-0.41, 0.83]
Scarano ■ 2020	36.9	4.2	20	35	2.8	20	21.0%	0.52	[-0.11, 1.15]
<b>Total (95% CI)</b>			<b>93</b>			<b>93</b>	<b>100.0%</b>	<b>0.80</b>	<b>[0.23, 1.38]</b>


Heterogeneity: Tau<sup>2</sup> = 0.31; Chi<sup>2</sup> = 14.07, df = 4 (P = 0.007); I<sup>2</sup> = 72%  
Test for overall effect: Z = 2.72 (P = 0.006)



### Skin temperature below a surgical mask during use

Study or Subgroup	surgical mask			no mask			Weight	Year	Std. Mean Difference	
	Mean	SD	Total	Mean	SD	Total			IV, Random, 95% CI	95% CI
Roberge 2012	33.7	0.88	20	31.94	1.1	20	49.1%	2012	1.73	[0.99, 2.47]
Scarano 2020	35.9	3.4	20	35.2	3.1	20	50.9%	2020	0.21	[-0.41, 0.83]
<b>Total (95% CI)</b>			<b>40</b>			<b>40</b>	<b>100.0%</b>		<b>0.96</b>	<b>[-0.53, 2.45]</b>


Heterogeneity: Tau<sup>2</sup> = 1.04; Chi<sup>2</sup> = 9.55, df = 1 (P = 0.002); I<sup>2</sup> = 90%  
Test for overall effect: Z = 1.26 (P = 0.21)



### Skin temperature below a N95 mask during use

Study or Subgroup	N95 mask			no mask			Weight	Std. Mean Difference	
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	95% CI
Kim 2016	35	0.7	12	34.8	0.7	12	29.1%	0.28	[-0.53, 1.08]
Park 2020	35.133	1.229	21	33.5	1.235	21	34.5%	1.30	[0.63, 1.97]
Scarano 2020	36.9	4.2	20	35	2.8	20	36.4%	0.52	[-0.11, 1.15]
<b>Total (95% CI)</b>			<b>53</b>			<b>53</b>	<b>100.0%</b>	<b>0.72</b>	<b>[0.12, 1.32]</b>

Heterogeneity: Tau<sup>2</sup> = 0.15; Chi<sup>2</sup> = 4.41, df = 2 (P = 0.11); I<sup>2</sup> = 55%  
Test for overall effect: Z = 2.35 (P = 0.02)



## B) Humidity (air humidity in % under mask)

### Humidity of breathing air with face mask compared to no mask

Study or Subgroup	mask			no mask			Weight	Year	Std. Mean Difference	
	Mean	SD	Total	Mean	SD	Total			IV, Random, 95% CI	95% CI
Roberge 2012	91.49	8.8	20	53.19	17.66	20	52.4%	2012	2.69	[1.81, 3.57]
Kim 2016	82.8	16.6	12	56	12.8	12	47.6%	2016	1.75	[0.78, 2.71]
<b>Total (95% CI)</b>			<b>32</b>			<b>32</b>	<b>100.0%</b>		<b>2.24</b>	<b>[1.32, 3.17]</b>

Heterogeneity: Tau<sup>2</sup> = 0.22; Chi<sup>2</sup> = 2.02, df = 1 (P = 0.16); I<sup>2</sup> = 50%  
Test for overall effect: Z = 4.75 (P < 0.00001)





Figure 5

Forest (left) and Funnel plots (right) of meta-analysis of physical outcomes while wearing a face mask. **A** shows results for temperature of skin, **B** for air humidity underneath the face mask. All mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If studies examine two different mask types in parallel, the corresponding studies are marked: □ =surgical mask ■ =N95 mask.

**A:** Skin covered by mask has a significantly higher temperature during rest and activity. This could be found for general mask use and for N95 mask use but not for surgical mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded higher temperatures than the surgical mask, but this could be analysed further due to lack of further studies comparing both conditions.

**B:** The dead space covered by mask has a significantly higher air humidity during rest and activity.



**Figure 6**

Forest and Funnel plots of meta-analysis of measured discomfort (A), itch (B), exertion (C), shortness of breath (D), perceived heat (E) and humidity (F) during face mask use (VAS, Likert-scales or similar). All face mask types are initially considered together, later subgroups (surgical and N95) are evaluated. If

studies examine two different mask types in parallel, the corresponding studies are marked: =surgical mask =N95 mask.

**A:** Perceived discomfort is significantly higher in face mask use in the pooled analysis. This could be found for general mask use, in the subgroup analysis for surgical-, and for N95 mask use. A pooled analysis comparing both conditions (surgical mask and N95 mask) resulted in statistically significant higher discomfort rates for the N95 mask than the surgical mask.

**B:** An overall significance for itching could be found for mask use. Also in N95 mask use the perceived itching was statistically significantly elevated according to the pooled subgroup analysis.

**C:** Perceived exertion is significantly higher in mask use. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher exertion rates than the surgical masks.

**D:** Perceived shortness of breath is significantly higher in mask use in the pooled analysis.



**E:** Perceived heat is significantly higher in the pooled analysis.

**F:** Perceived humidity is significantly higher in mask use. The subgroup analysis revealed a statistical significance for an increase in humidity perception using a surgical mask. In studies evaluating both conditions (surgical and N95 mask) the N95 mask yielded always higher humidity perception rates than the surgical mask. A pooled analysis resulted in a statistical significance for higher humidity perception in N95 masks than surgical masks.



# Meta-analysis of N95 mask vs surgical mask

## A) Biochemical comparison

### Blood oxygen saturation (SpO<sub>2</sub>) when using a N95 mask vs surgical mask





### Carbon dioxide blood content (mmHg) in N95 mask use vs surgical mask





## B) Cardiorespiratory comparison



### Ventilation (l/min) when using a N95 mask vs surgical mask



### Heart rate (beats/min) when using a N95 mask vs surgical mask





### Systolic blood pressure (mmHg) when using a N95 vs surgical mask





## C) Measured subjective sensations comparison

### Perceived discomfort when using a N95 mask vs surgical mask



### Perceived exertion when using a N95 mask vs surgical mask



### Perceived humidity when using a N95 mask vs surgical mask






Figure 7

Results comparing the N95 to the surgical mask in the meta-analysis. Forest (left) and Funnel plots (right) of meta-analysis of diverse outcomes while wearing a N95 mask vs surgical mask are shown. **A** depicts the biochemical, **B** the cardiorespiratory outcomes and **C** the subjective sensations outcomes.


N95 mask leads to measurably less favourable results compared to the surgical mask, significantly for oxygenation (decrease), heart rate (increase), discomfort and humidity (both increases). This trend was also evident for minute volume (decrease), CO<sub>2</sub> and systolic blood pressure (both increases), but in those comparisons not statistically significant due to too few includable studies.

## Meta-analysis of pooled symptom prevalence while wearing a face mask


### Headache prevalence when using a face mask




### Acne prevalence when using a face mask




### Skin irritation prevalence when using a face mask




### Shortness of breath prevalence when using a face mask




### Heat prevalence when using a face mask




### Itch prevalence when using a face mask



### Voice disorder prevalence when using a face mask




### Dizziness prevalence when using a face mask



**Figure 8**

Forest (left) and Funnel plots (right) of meta-analysis of pooled symptom prevalence while wearing a face mask. Headache (62%), acne (38%), skin irritation (36%), shortness of breath (33%), heat (26%), itch (26%), voice disorder (23%) and dizziness (5%) while wearing a mask are significant in the evaluated population (n=8128).

**Metaanalytically measured biochemical and physical effects of face masks**



**Figure 9**

Summary of pooled metaanalytic evaluation of biochemical (A) and physical effects (B) during face mask use. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the confidence intervals.

**A:** For carbon dioxide rise in the blood there is a medium effect size of >0.5 and for oxygen drop a small effect size of >0.2 regarding the standard mean difference values thresholds according to Cohen 1988.

**B:** For elevated Humidity and Temperature rise under the face mask there is a strong effect size of  $\geq 0.8$ .

The metaanalytical statistical data were as follows:

Oxygen (SpO<sub>2</sub>): SMD -0.24, 95% CI -0.38 to -0.11, Z=3.53, p=0.0004;

Carbon dioxide (PtCO<sub>2</sub>, ETCO<sub>2</sub>, PaCO<sub>2</sub>): SMD +0.64, 95% CI 0.31 to 0.96, Z=3.86, p=0.0001;

Humidity: SMD +2.24, 95% CI 1.32 to 3.17), Z=4.75, p<0.00001;

Temperature: SMD +0.8, 95% CI 0.23 to 1.38, Z=2.72, p=0.008.

## Metaanalytically measured cardiorespiratory effects of face masks




Figure 10

Summary of pooled metaanalytic evaluation of cardiorespiratory effects during face mask use. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the

confidence intervals.

Clear effects for a decrease in ventilation and tidal volume are illustrated, no effect for respiratory rate and weak to low effect for increase in heart rate and systolic blood pressure. For ventilation there is a medium effect size of  $>0.5$  with a small effect size of  $>0.2$  for tidal volume of the standard mean difference values according to Cohen 1988.

The meta-analytical statistical data were as follows:

Ventilation: SMD -0.72, 95% CI -0.99 to -0.46,  $Z=5.36$ ,  $p<0.00001$ ;


Tidal volume: SMD -0.37, 95% CI -0.63 to -0.11,  $Z=2.82$ ,  $p=0.005$ ;

Respiratory rate: SMD +0.01, 95% CI -0.29 to 0.30,  $Z=0.08$ ,  $p=0.94$ ;

Heart rate: SMD +0.11, 95% CI -0.05 to 0.28,  $Z=1.34$ ,  $p=0.18$ ;

Systolic blood pressure: SMD +0.17, 95% CI 0.03 to 0.32,  $Z=2.39$ ,  $p=0.02$ .

## Trial duration in studies investigating predominantly physiological, chemical and physical mask effects




## Figure 11

Illustration of the duration of studies in which measurements were made on mask effects (physical, biochemical, and physiological) in 934 participants.

The median is 18 minutes (yellow dotted line) with an interquartile range of 50 . The study with the longest experimental duration included 21 subjects, corresponding to 2.2% of the total population studied.

Striking not only is a very short trial time compared to the everyday scenarios workday and school attendance (see interrupted, auxiliary lines in blue and red), but also a strongly deviating mask exposure duration with outliers (mean of 45.8 minutes with standard deviation of 69.9). Therefore, the mean is not an appropriate parameter to characterize this distribution.

## Metaanalytically measured symptoms while wearing face masks



**Figure 12**

Summary of pooled metaanalytic evaluation of face mask-wearing sensations measured with standardised Borg-, Likert-, VAS-scales or similar. The height of the bars reflects the SMD (standard mean difference), their error bars correspond to the confidence intervals.

Five out of 6 complaint categories (83%) are above the strong effect size threshold of >0.8 of the standard mean difference values according to Cohen 1988.

The metaanalytical statistical data were as follows (SMD=standard mean difference):

Itch: SMD +1.57, 95%CI -0.08 to 3.23, Z=1.86, p=0.06;

Shortness of breath: SMD +1.46, 95% CI 0.42 to 2.50, Z=2.75, p=0.006;

Discomfort: SMD +1.16, 95% CI 0.58 to 1.73, Z=3.94, p<0.0001;

Exertion: SMD +0.9, 95% CI 0.57 to 1.23, Z=5.31, p<0.00001;

Humidity: SMD +0.9, 95% CI 0.34 to 1.46, Z=3.17, p=0.002;

Heat: SMD +0.77, 95% CI 0.29 to 1.26, Z=3.11, p=0.002.

## Metaanalytically pooled prevalence of symptoms with face mask




Figure 13

Representation of symptom prevalence in % during face mask use as the area of the circles. Along the X-axis, the main recorded symptoms are listed. The higher the prevalence, the bigger the circles and the more often the symptoms.

The Y-axis gives the probability of non-random occurrence of the symptoms and includes the statistical Z-value. Thus, the higher the circles are arranged, the more robust is the relationship to face mask wearing.

The meatanalytical statistical data were as follows:

Headache: 62% (95% CI 48-75%), Z=8.77, p<0.00001;

Acne: 38% (95% CI 22-54%), Z=4.58, p<0.00001:

Skin irritation: 36% (95% CI 24-49%), Z=5.61, p<0.00001;

Shortness of breath: 33% (95% CI 23-44%), Z=6.28, p<0.00001;

Heat: 28% (95% CI 15-0.37%), Z=4.72, p<0.00001;

Itch: 26% (95% CI 15-36%), Z=4.77, p<0.00001;

Voice disorder 23% (95% CI 2-43%), Z=2.15, p<0.03;

Dizziness 5% (95% CI 1-9%), Z=2.5, p=0.01.